FINITE ELEMENT EXTERIOR CALCULUS FOR EVOLUTION PROBLEMS

ANDREW GILLETTE, MICHAEL HOLST, AND YUNRONG ZHU

ABSTRACT. Arnold, Falk, and Winther [Bull. Amer. Math. Soc. 47 (2010), 281–354] showed that mixed variational problems, and their numerical approximation by mixed methods, could be most completely understood using the ideas and tools of *Hilbert com*plexes. This led to the development of the Finite Element Exterior Calculus (FEEC) for a large class of linear elliptic problems. More recently, Holst and Stern [Found. Comp. Math. 12:3 (2012), 263–293 and 363–387] extended the FEEC framework to semi-linear problems, and to problems containing variational crimes, allowing for the analysis and numerical approximation of linear and nonlinear geometric elliptic partial differential equations on Riemannian manifolds of arbitrary spatial dimension, generalizing surface finite element approximation theory. In this article, we develop another distinct extension to the FEEC, namely to parabolic and hyperbolic evolution systems, allowing for the treatment of geometric and other evolution problems. Our approach is to combine the recent work on the FEEC for elliptic problems with a classical approach to solving evolution problems via semi-discrete finite element methods, by viewing solutions to the evolution problem as lying in time-parameterized Hilbert spaces (or Bochner spaces). Building on classical approaches by Thomée for parabolic problems and Geveci for hyperbolic problems, we establish a priori error estimates for Galerkin FEM approximation in the natural parametrized Hilbert space norms. In particular, we recover the results of Thomée and Geveci for two-dimensional domains and lowest-order mixed methods as special cases, effectively extending their results to arbitrary spatial dimension and to an entire family of mixed methods. We also show how the Holst and Stern framework allows for extensions of these results to certain semi-linear evolution problems.

CONTENTS

1.	Introduction	2
2.	Semi-Discrete FEM Error Estimates for Evolution Problems	4
3.	The Finite Element Exterior Calculus	7
4.	A Priori Error Estimates for Parabolic Problems	9
5.	A Priori Error Estimates for Hyperbolic Problems	14
6.	Semi-linear Evolution Problems	18
7.	Concluding Remarks	22
Appendix A. Explanation of Semi-Linear Error Estimates		23
References		24

Date: October 16, 2018.

Key words and phrases. FEEC, elliptic equations, evolution equations, nonlinear equations, approximation theory, nonlinear approximation, inf-sup conditions, *a priori* estimates.

1. INTRODUCTION

More than two decades of research on linear mixed variational problems, and their numerical approximation by mixed methods, recently culminated in the seminal work of Arnold, Falk, and Winther [3]. The authors show how such problems are most completely understood using the ideas and tools of *Hilbert complexes*, leading to the development of the Finite Element Exterior Calculus (FEEC) for elliptic problems. In two related articles [20, 21], Holst and Stern extended the Arnold–Falk–Winther framework to include *variational crimes*, allowing for the analysis and numerical approximation of linear and nonlinear geometric elliptic partial differential equations on Riemannian manifolds of arbitrary spatial dimension, generalizing the existing surface finite element approximation theory in several directions.

In this article, we extend the FEEC in another direction, namely to parabolic and hyperbolic evolution systems. Our approach is to combine the recent work on the FEEC for elliptic problems with a classical approach to solving evolution problems using semidiscrete finite element methods, by viewing solutions to the evolution problem as lying in time-parameterized Banach (or *Bochner*) spaces. Building on classical approaches by Thomée for parabolic problems and Geveci for hyperbolic problems, we establish *a priori* error estimates for Galerkin FEM approximation in the natural parametrized Hilbert space norms. In particular, we recover the results of Thomée and Geveci for twodimensional domains and the lowest-order mixed method as a special case, effectively extending their results to arbitrary spatial dimension and to an entire family of mixed methods. We also show how the Holst and Stern framework allows for extensions of these results to certain semi-linear evolution problems.

To understand why the finite element exterior calculus (FEEC) has emerged in a natural way to become a major mathematical tool in the development of numerical methods for PDE, we recall one of the many examples presented at length in [3]. Consider the vector Laplacian over a domain $\Omega \subset \mathbb{R}^3$:

$$-\Delta u := -$$
 grad div $u +$ curl curl u ,

with the boundary conditions $u \cdot n = 0$ and curl $u \times n = 0$ on $\partial\Omega$. Given an L^2 vector field data f, the natural variational formulation of the problem is: Find $u \in H(\text{curl}; \Omega) \cap H_0(\text{div}; \Omega)$ such that

$$\int_{\Omega} \left[(\nabla \cdot u) (\nabla \cdot v) + (\nabla \times u) \cdot (\nabla \times v) \right] dx = \int_{\Omega} f \cdot v \, dx, \, \forall v \in H(\operatorname{curl}; \Omega) \cap H_0(\operatorname{div}; \Omega).$$
(1.1)

By introducing an intermediate variable $\sigma := -\text{div } u$, a natural alternative formulation is the following *mixed* form: find $(\sigma, u) \in H^1(\Omega) \times H(\text{curl}; \Omega)$ such that

$$\int_{\Omega} (\sigma \tau - u \cdot \nabla \tau) \, dx = 0, \quad \forall \tau \in H^1(\Omega), \tag{1.2}$$

$$\int_{\Omega} [\nabla \sigma \cdot v + (\nabla \times u) \cdot (\nabla \times v)] \, dx = \int_{\Omega} f \cdot v \, dx, \quad \forall v \in H(\operatorname{curl}; \Omega).$$
(1.3)

Using the standard finite element approach based on the non-mixed formulation (1.1) (e.g. using continuous piecewise linear vector functions) can yield incorrect results if the domain has certain geometric features (e.g. domains with re-entrant corners) or topological features (e.g. non-simply connected domains). A standard finite element approach based on the mixed formulation (1.2)-(1.3), on the other hand, suffers neither of these difficulties and typically works extremely well. The explanation for why one approach fails and the other succeeds lies in the fundamental mathematical structures underlying the finite element method. In case of the domain with re-entrant corners, the failure of the non-mixed approach is due to the nondensity of $H^1(\Omega)$ in $H(\operatorname{curl}) \cap H_0(\operatorname{div})$, as discussed in [19, Remark 19] and [3, Section 2.3]. This non-density leads to a problem of *inconsistency*, i.e. that the discrete approximation of the operators and data do not approximate the continuous problem correctly as the mesh size is taken to zero. On the other hand, the error due to topological features, such as non-simply connected domains, can be traced to the presence of non-zero *harmonic vector fields* on the domain, i.e. vector fields that are both curl-free and divergencefree. The mixed formulation turns out to be both consistent and respectful of non-zero harmonic vector fields while the standard formulation does not. A natural question is then: What is an appropriate mathematical framework for understanding these problems abstractly so that a methodical construction of "good" finite element methods can be carried out for these and similar initial boundary value problems?

The answer turns out to be *Hilbert Complexes*. Hilbert complexes were originally studied in [8] as a way to generalize certain properties of elliptic complexes, particularly the Hodge decomposition and other aspects of Hodge theory. A *Hilbert complex* (W, d) consists of a sequence of Hilbert spaces W^k , along with closed, densely-defined linear maps $d^k : V^k \subset W^k \to V^{k+1} \subset W^{k+1}$, possibly unbounded, such that $d^k \circ d^{k-1} = 0$ for each k.

$$\cdots \longrightarrow V^{k-1} \xrightarrow{\mathrm{d}^{k-1}} V^k \xrightarrow{\mathrm{d}^k} V^{k+1} \longrightarrow \cdots$$

This Hilbert complex is said to be *bounded* if d^k is a bounded linear map from W^k to W^{k+1} for each k, i.e., (W, d) is a cochain complex in the category of Hilbert spaces. It is said to be *closed* if the image $d^k V^k$ is closed in W^{k+1} for each k. It was shown in [2, 3] that Hilbert complexes are also a convenient abstract setting for mixed variational problems and their numerical approximation by mixed finite element methods, providing the foundation of a framework called *finite element exterior calculus*. This line of research is the culmination of several decades of work on mixed finite element methods and computational electromagnetics [6, 17, 25, 26]. The most important example of a Hilbert complex for our purposes of the FEEC arises from the de Rham complex of smooth differential forms on a domain or manifold.

The main developments in FEEC to date have been for linear (and now semi-linear) elliptic problems such as Poisson's equation

$$-\Delta u = f.$$

Our goal here is to expand the scope of this analysis to include parabolic linear (and semi-linear) equations such as the heat equation,

$$(\partial_t - \Delta)u = f$$

and hyperbolic equations such as the wave equation,

$$(\partial_{tt} - \Delta)u = f.$$

The exterior calculus framework treats Δ as $(d + \delta)^2$, where d is the exterior derivative operator and δ its adjoint. The incorporation of the time derivative operation ∂_t into this framework, however, has not been previously considered. To remedy this, we develop the most natural extension of FEEC theory to evolution problems: a generalization of the semi-discrete method often called the 'method of lines.' This approach involves the discretization of the spatial part of the differential operator, leaving the time variable continuous. It can be viewed as introducing a time parameter into the discrete (Hilbert) spaces that have been developed for elliptic problems. The FEEC terminology provides a clear and consistent notation for bounding errors in mixed methods accumulated over a finite time interval.

We note that there is another approach to solving evolution problems with finite elements, namely using a complete discretization of space-time. This tactic allows for the dynamical change of the underlying discrete approximation spaces in both space and time. Such an approach gives rise to space-time adaptivity, and is potentially the most flexible and powerful approach to the numerical treatment of parabolic and hyperbolic evolution problems. This approach, which we will consider in a second article, is most naturally formulated using geometric calculus, a well-studied mathematical structure for time-dependent problems. In the current article, we focus on extending FEEC to semidiscrete methods using time-parameterized Banach (i.e. Bochner) space norm estimates for the method-of-lines approach.

Finally, we note that the work presented here was developed simultaneously and independently from a related project by Arnold and Chen [1] for generalized Hodge-Laplacian style linear parabolic problems. Our focus in this work is to extend the *scalar* Hodge-Laplacian to both linear and semi-linear parabolic problems as well as linear hyperbolic problems, as this touches the existing literature on semi-discrete methods in the broadest fashion. The pairing of these two results will lead to further insight in a variety of research directions.

Outline and summary of contributions. The remainder of the paper is structured as follows. In Section 2, we review the classical semi-discrete mixed finite element method error estimates for parabolic problems (due to Thomée [33] and others) and for hyperbolic problems (due to Geveci [15] and others). In Section 3, we give a very brief overview of the Finite Element Exterior Calculus and recall some relevant results. In Section 4, we combine the classical approach to semi-discrete methods with modern FEEC theory to establish some basic a priori error estimates for Galerkin mixed finite element methods for parabolic problems. The main result is Theorem 4.2, which exploits the FEEC framework to obtain a classification of spatial finite element spaces that give optimal order convergence rates in Bochner norms. In Section 5, we carry out a similar analysis for hyperbolic problems, resulting in the error estimate given in Theorem 5.2, a simultaneous sharpening of the result by Geveci for problems in two dimensional domains and a generalization to problems on n dimensional domains. Our results recover the estimates of Thomée and Geveci for two-dimensional domains and the lowest-order mixed method as a special case, effectively extending their results to arbitrary spatial dimension and to an entire family of mixed methods. In Section 6, we employ the results of Holst and Stern [21] to extend our parabolic estimates to a class of semi-linear evolution PDE, as encapsulated by Theorem 6.2. Finally, in Section 7, we draw conclusions and make remarks on future directions. The appendices contain some technical details relating to the application of the semi-linear error estimates in the case k = n and a review of some standard existence and uniqueness results for abstract parabolic and hyperbolic problems using the terminology of Bochner spaces.

2. SEMI-DISCRETE FEM ERROR ESTIMATES FOR EVOLUTION PROBLEMS

We begin by reviewing semi-discrete finite element methods and their *a priori* error estimates for parabolic and hyperbolic PDE systems. We focus in each case on a relatively simple, well-studied system of interest to modeling communities, namely, the heat equation (parabolic) and the wave equation (hyperbolic). To avoid complicating the problem statements, we assume the given data f, g, u_0 and u_1 to be as regular as necessary for the problem context, unless otherwise specified. The **heat equation** is: find u(x, t) such that

$$u_t - \Delta u = f \quad \text{in } \Omega, \quad \text{for } t > 0$$

$$u = 0 \quad \text{on } \partial\Omega, \quad \text{for } t > 0 \quad \text{with } u(\cdot, 0) = g \text{ in } \Omega.$$
(2.1)

We review the approach to Galerkin methods for this problem as presented in Thomée [33] for domains $\Omega \subset \mathbb{R}^2$. His approach is based on work with Johnson [22] and builds upon prior analysis of elliptic projection [7]. A similar approach, restricted to $\Omega \subset \mathbb{R}^2$, was carried out by Garcia in [14]. Let $\sigma = \nabla u$ and define the mixed, weak form problem: Given f and g, find $(u(t), \sigma(t)) \in L^2(\Omega) \times H(\operatorname{div}; \Omega)$ such that

$$\begin{aligned} &(u_t,\phi) - (\operatorname{div}\sigma,\phi) &= (f,\phi), \quad \forall \ \phi \in L^2(\Omega), \quad t > 0 \\ &(u,\operatorname{div}\omega) + (\sigma,\omega) &= 0, \quad \forall \ \omega \in H(\operatorname{div};\Omega), \quad t > 0, \quad u(0) = g. \end{aligned}$$
 (2.2)

The semi-discrete problem is then to find $(u_h(t), \sigma_h(t)) \in S_h \times H_h \subset L^2(\Omega) \times H(\operatorname{div}; \Omega)$ such that

$$\begin{aligned} & (u_{h,t},\phi_h) - (\operatorname{div}\sigma_h,\phi_h) &= (f,\phi_h), \quad \forall \ \phi_h \in S_h, \quad t > 0 \\ & (u_h,\operatorname{div}\omega_h) + (\sigma_h,\omega_h) &= 0, \qquad \forall \ \omega_h \in H_h, \quad t > 0, \quad u_h(0) = g_h. \end{aligned}$$

$$(2.3)$$

where g_h is an approximation of g in S_h . With bases for S_h and H_h , the matrix form of the discrete problem is

$$\begin{split} &AU_t - B\Sigma = F,\\ &B^TU + D\Sigma = 0, \ \ \text{for} \ t > 0, \ \ U(0) \ \text{given}, \end{split}$$

where U and Σ are vectors corresponding to u_h and σ_h . It is easily seen that the matrices A and D are positive definite. Eliminating Σ , we have the system of ODEs

$$AU_t + BD^{-1}B^T U = F$$
, for $t > 0$, $U(0)$ given,

which by standard results in ODE theory has a unique solution.

Thomée uses discontinuous linear elements for S_h and piecewise quadratic elements for H_h . He defines the solution operator $T_h : L^2 \to S_h$ given by $T_h f = u_h$ for the corresponding elliptic problem and sets

$$g_h := R_h g := -T_h \Delta g.$$

For $g_h = R_h g$ and $t \ge 0$, Thomée derives the estimates

$$||u_h(t) - u(t)||_{L^2} \le ch^2 \left(||u(t)||_{H^2} + \int_0^t ||u_t||_{H^2} \, ds \right), \tag{2.4}$$

$$||\sigma_h(t) - \sigma(t)||_{L^2} \le ch^2 \left(||u(t)||_{H^3} + \left(\int_0^t ||u_t||_{H^2}^2 \, ds \right)^{1/2} \right). \tag{2.5}$$

Note that these estimates are for a fixed time value t and restricted to a particular choice of finite elements in 2D.

We now turn to the **wave equation**: find u(x, t) such that

$$u_{tt} - \Delta u = f \quad \text{in } \Omega, \quad \text{for } t > 0,$$

$$u = 0 \quad \text{on } \partial \Omega, \quad \text{for } t > 0 \quad \text{with } u(\cdot, 0) = u_0 \text{ in } \Omega,$$

and $u_t(\cdot, 0) = u_1 \text{ in } \Omega$
(2.6)

There are two approaches to defining a mixed weak form of this problem. The first is very similar to the parabolic case: find $(u(t), \sigma(t)) \in L^2(\Omega) \times H(\text{div}; \Omega)$ such that

$$(u_{tt}, \phi) - (\operatorname{div} \sigma, \phi) = (f, \phi), \quad \forall \phi \in L^{2}(\Omega), \quad t > 0,$$

$$(u, \operatorname{div} \omega) + (\sigma, \omega) = 0, \quad \forall \omega \in H(\operatorname{div}; \Omega), \quad t > 0,$$

$$u(0) = u_{0},$$

$$u_{t}(0) = u_{1}.$$
(2.7)

It is difficult to derive estimates for the numerical approximation of (2.7) akin to those found in the parabolic case due to the second derivatives appearing in the formulation. Some attempts at estimates along these lines for $\Omega \subset \mathbb{R}^2$ have been given by Baker [4] and Cowsar, Dupont and Wheeler [10, 11].

For the purpose of extending the FEEC framework, we find the 'velocity-stress' formulation of the problem and the results of Geveci [15] to be more useful. This formulation solves for $\mu := u_t$ instead of u: Given f, u_0 , and u_1 , find $(\mu(t), \sigma(t)) \in L^2(\Omega) \times H(\text{div}; \Omega)$ such that

$$\begin{aligned} (\mu_t, \phi) - (\operatorname{div} \sigma, \phi) &= (f, \phi), &\forall \phi \in L^2(\Omega), \quad t > 0, \\ (\mu, \operatorname{div} \omega) + (\sigma_t, \omega) &= 0, \quad \forall \omega \in H(\operatorname{div}; \Omega), \quad t > 0, \\ \mu(0) &= u_1, \\ \sigma(0) &= \nabla u_0. \end{aligned}$$

$$(2.8)$$

The semi-discrete problem is then to find $(\mu_h(t), \sigma_h(t)) \in S_h \times H_h$ such that

$$\begin{aligned} (\mu_{h,t},\phi_h) - (\operatorname{div} \sigma_h,\phi_h) &= (f,\phi_h), & \forall \phi_h \in S_h, \quad t > 0, \\ (\mu_h,\operatorname{div} \omega_h) + (\sigma_{h,t},\omega_h) &= 0, & \forall \omega_h \in H_h, \quad t > 0, \\ \mu_h(0) &= u_{1,h}, \\ \sigma_h(0) &= (\nabla u_0)_h, \end{aligned}$$

$$(2.9)$$

where $u_{1,h}$ is an approximation of u_1 in S_h and $(\nabla u_0)_h$ is an approximation of ∇u_0 . Again, bases for S_h and H_h reduce the discrete problem to a matrix formulation:

$$AW_t - B\Sigma = F,$$

 $B^TW + D\Sigma_t = 0, \text{ for } t > 0, W(0), \Sigma(0) \text{ given},$

where W and Σ are vectors corresponding to μ_h and σ_h and A and D are symmetric, positive definite matrices. As Geveci [15, p. 248] explains, this can be reduced to a single iterative system of the form

$$(D+k^2B^TA^{-1}B)\Sigma^{n+1} = G,$$

where k denotes the time step in an implicit Euler time-differencing scheme.

To derive an error estimate for the velocity-stress discretization, Geveci states the need for projection operators from H(div) to H_h and from L^2 to S_h satisfying certain approximation properties. He explains that such operators exists for a variety of finite element spaces in \mathbb{R}^2 , e.g. the Raviart-Thomas spaces [28], allowing the following result. For

$$1 \leq s \leq r \text{ with } r \geq 2,$$

$$||\mu_{h}(t) - \mu(t)||_{L^{2}} + ||\sigma_{h}(t) - \sigma(t)||_{L^{2}} \leq c \left(||u_{1} - u_{1,h}||_{L^{2}} + ||\nabla u_{0} - (\nabla u_{0})_{h}||_{L^{2}}\right) + ch^{s} \left(||u_{1}||_{s} + ||\nabla u_{0}||_{s} + \int_{0}^{t} (||\mu_{t}(\tau)||_{s} + ||\sigma_{t}(\tau)||_{s} d\tau\right).$$

$$(2.10)$$

Like estimates (2.4) and (2.5) for the parabolic problem, (2.10) says that the approximation error can be controlled in L^2 norm at any time t by the H^s norm of the initial conditions plus the accumulated norm of the variables up to time t. It is these types of estimates that the FEEC framework can refine, simplify, and generalize to arbitrary spatial dimension n.

3. THE FINITE ELEMENT EXTERIOR CALCULUS

The finite element exterior calculus (FEEC) provides an elegant mathematical framework for deriving error estimates for a large class of elliptic PDE. We now give a brief overview of the notation and certain main results from FEEC which are relevant to this paper. We refer the reader to the seminal papers of Arnold, Falk, and Winther [2, 3] for additional explanation.

Let Ω be a bounded *n*-manifold embedded in \mathbb{R}^n and assume Ω has a piecewise smooth, Lipschitz boundary. The space of L^2 -bounded continuous differential k-forms on Ω is given by

$$L^{2}\Lambda^{k}(\Omega) := \left\{ \sum_{i} a_{i} dx_{i} \in \Lambda^{k}(\Omega) : a_{i} \in L^{2}(\Omega) \quad \forall i \right\},\$$

where *i* ranges over all strictly increasing sequences of *k* indices chosen from $\{1, \ldots, n\}$. The exterior derivative operator $d_k : \Lambda^k(\Omega) \to \Lambda^{k+1}(\Omega)$ acts on these spaces to form a Hilbert complex $(L^2\Lambda, d)$. The associated domain complex is the sequence of spaces $H\Lambda^k := \text{domain}(d_k) \subset L^2\Lambda^k(\Omega)$, commonly called the L^2 deRham complex:

$$0 \longrightarrow H\Lambda^0 \xrightarrow{d_0} H\Lambda^1 \xrightarrow{d_1} \cdots \xrightarrow{d_{n-1}} H\Lambda^n \longrightarrow 0.$$

The norm on each space is the graph norm associated to d, i.e.

$$(u,v)_{H^k(\Omega)} := (u,v)_{\Lambda^k(\Omega)} + (d_k u, d_k v)_{\Lambda^{k+1}(\Omega)}$$

We note that in any dimension n, the beginning and end of the L^2 deRham complex can be understood in terms of traditional Sobolev spaces and differential operators:

$$0 \longrightarrow H^1(\Omega) \xrightarrow{\operatorname{grad}} \cdots \xrightarrow{\operatorname{div}} L^2(\Omega) \longrightarrow 0$$

A major conclusion of FEEC is that stable finite element methods for elliptic PDE must seek solutions in finite dimensional subspaces $\Lambda_h^k \subset H\Lambda^k$ that satisfy certain key approximation properties. First, the subspaces should form a subcomplex of the L^2 deR-ham complex, meaning $d\Lambda_h^k \subset \Lambda_h^{k+1}$. Second, Λ_h^k should have sufficient approximation that upper bounds on $\inf_{v \in \Lambda_h^k} ||u - v||_{H\Lambda^k}$ can be ensured for some or all $u \in H\Lambda^k$. Third, there must exist bounded cochain projections $\pi_h^k : H\Lambda^k \to \Lambda_h^k$ which are invariant on Λ_h^k , commute with the exterior derivative operators, and provide a bound $||\pi_h^k v||_{H\Lambda^k} \leq c ||v||_{H\Lambda^k}$ for all $v \in H\Lambda^k$.

In the context of the deRham complex, all these properties are shown to be provided for by two canonical classes of piecewise degree r polynomials associated to a simplicial mesh \mathcal{T} of Ω . Let \mathcal{P}_r denote polynomials in n variables of degree at most r and $\mathcal{H}_r \subset \mathcal{P}_r$ the subspace of homogeneous polynomials. The first class, denoted $\mathcal{P}_r\Lambda^k(\mathcal{T})$, consists of all k-forms with coefficients belonging to \mathcal{P}_r on each n-simplex of \mathcal{T} . The second class, denoted $\mathcal{P}_r^-\Lambda^k(\mathcal{T})$, interleaves with the first class, i.e.

$$\mathcal{P}_{r-1}\Lambda^k(\mathcal{T}) \subsetneq \mathcal{P}_r^-\Lambda^k(\mathcal{T}) \subsetneq \mathcal{P}_r\Lambda^k(\mathcal{T}).$$

To define $\mathcal{P}_r^-\Lambda^k(\mathcal{T})$, first define X be the vector field on \mathbb{R}^n such that X(x) is the vector based at $x \in \mathbb{R}^n$ that points opposite to the origin with length |x|. Define $\mathcal{P}_r^-\Lambda^k(\mathcal{T}) :=$ $\mathcal{P}_r\Lambda^k \oplus \kappa \mathcal{H}_{r-1}\Lambda^{k+1}$, a direct sum, where κ is defined by contraction with X. The map κ is called the *Koszul differential* and gives rise to the *Koszul complex*. This is elaborated upon in detail in the work of Arnold, Falk and Winther e.g. [3, p. 328].

For n = 3, we have the following correspondences between the FEEC notation of finite element spaces and traditional finite element spaces.

$$\mathcal{P}_{r+1}\Lambda^2(\mathcal{T}) = \text{Nédélec 2nd-kind } H(\text{div}) \text{ elements of degree} \le r+1 \text{ (see [26])}$$

$$\mathcal{P}_{r+1}^{-}\Lambda^2(\mathcal{T}) = \text{Nédélec 1st-kind } H(\text{div}) \text{ elements of order } r+1 \text{ (see [25])}$$

 $\mathcal{P}_{r+1}^{-}\Lambda^{3}(\mathcal{T}) = \mathcal{P}_{r}\Lambda^{3}(\mathcal{T}) = \text{discontinuous elements of degree} \leq r$

Hence, in the case of the deRham complex, FEEC recovers well-known finite element spaces while at the same time describing their generalization to arbitrary spatial dimensions.

The last piece of FEEC used in this work is the existence of smoothed projection operators

$$\pi_h^k : L^2 \Lambda^k \to L^2 \Lambda_h^k \quad \text{where } \Lambda_h^k \in \{ \mathcal{P}_r \Lambda^k(\mathcal{T}), \ \mathcal{P}_r^- \Lambda^k(\mathcal{T}) \}.$$
(3.1)

These operators are shown, by virtue of their construction, to be uniformly bounded (in $L^2\Lambda^k$) with respect to h. An explicit construction of these operators can be found in the papers of Arnold, Falk, and Winther [2, 3]; the following theorem asserts some of their key properties.

Theorem 3.1 ([3] Theorem 5.9).

(i.) Let Λ_h^k be one of the spaces $\mathcal{P}_{r+1}^-\Lambda^k(\mathcal{T})$ or, if $r \geq 1$, $\mathcal{P}_r\Lambda^k(\mathcal{T})$. Then π_h^k is a projection onto Λ_h^k and satisfies

$$\left|\left|\omega - \pi_{h}^{k}\omega\right|\right|_{L^{2}\Lambda^{k}(\Omega)} \leq ch^{s} \left|\left|\omega\right|\right|_{H^{s}\Lambda^{k}(\Omega)}, \quad \omega \in H^{s}\Lambda^{k}(\Omega),$$

for $0 \leq s \leq r+1$. Moreover, for all $\omega \in L^2\Lambda^k(\Omega)$, $\pi_h^k \omega \to \omega$ in L^2 as $h \to 0$. (ii.) Let Λ_h^k be one of the spaces $\mathcal{P}_r\Lambda^k(\mathcal{T})$ or $\mathcal{P}_r^-\Lambda^k(\mathcal{T})$ with $r \geq 1$. Then

$$\left|\left|d(\omega - \pi_h^k \omega)\right|\right|_{L^2 \Lambda^k(\Omega)} \le ch^s \left|\left|d\omega\right|\right|_{H^s \Lambda^k(\Omega)}, \quad \omega \in H^s \Lambda^k(\Omega)$$

for $0 \leq s \leq r$. (iii.) Let $\Lambda_h^{k-1} \in \{\mathcal{P}_{r+1}\Lambda^{k-1}(\mathcal{T}), \mathcal{P}_{r+1}^-\Lambda^{k-1}(\mathcal{T})\}$ and $\Lambda_h^k = \mathcal{P}_{r+1}^-\Lambda^k(\mathcal{T})$ or, if r > 0, $\mathcal{P}_r\Lambda^k(\mathcal{T})$. Then $d\pi_h^{k-1} = \pi_h^k d$.

To handle the setting of parabolic and hyperbolic PDE, we now merge the notation of FEEC with the abstract framework of parametrized Banach spaces, also known as Bochner spaces. We follow prior approaches along these lines, especially [32, page 66] and [29]. Let X be a Banach space and I := (0, T) an interval with closure $I := \overline{I} = [0, T]$. Define the space

 $C(\mathring{I},X) := \{ u : \mathring{I} \to X \mid u \text{ bounded and continuous} \},$

and equip it with the norm

$$||u||_{C(\mathring{I},X)} := \sup_{t\in\mathring{I}} ||u(t)||_X.$$

The **Bochner space** $L^P(\mathring{I}, X)$ is then defined to be the completion of $C(\mathring{I}, X)$ with respect to the norm

$$||u||_{L^{p}(\mathring{I},X)} := \left(\int_{\mathring{I}} ||u(t)||_{X}^{p} dt\right)^{1/p}$$

The space $H^1(\mathring{I}, X)$ has an analogous norm

$$||u||_{H^{1}(\mathring{I},X)} := \left(\int_{\mathring{I}} ||u(t)||_{X}^{2} + \left\| \frac{d}{dt} u(t) \right\|_{X}^{2} dt \right)^{1/2}$$

We will commonly use $X = L^2 \Lambda^k$ or $X = H^s \Lambda^k$ where it is understood that the forms are defined over a spatial domain Ω . In the case of spatial-only norms, we will use the notation $||\cdot||_{L^2}$ or just $||\cdot||$ for the $L^2 \Lambda^k$ norm, and $||\cdot||_{H^s}$ for the $H^s \Lambda^k$ norm, where k will be clear from context.

4. A Priori Error Estimates for Parabolic Problems

We extend Thomée's error estimates from Section 2 to the broader class of elements and arbitrary spatial dimension allowed by FEEC with Bochner space norms. For simplicity, let $\Omega \subset \mathbb{R}^n$ be a contractible domain. Define the **mixed weak parabolic** problem: Given f, g, find $(u, \sigma) : I \to H\Lambda^n \times H\Lambda^{n-1}$ such that

$$\begin{aligned} (u_t,\phi) - (\operatorname{div} \sigma,\phi) &= (f,\phi), &\forall \phi \in H\Lambda^n, \quad t \in I, \\ (u,\operatorname{div} \omega) + (\sigma,\omega) &= 0, \quad \forall \omega \in H\Lambda^{n-1}, \quad t \in I, \\ u(0) &= g. \end{aligned}$$
 (4.1)

The semi-discrete parabolic problem is thus: Find $(u_h, \sigma_h) : I \to \Lambda_h^n \times \Lambda_h^{n-1}$ such that

$$(u_{h,t},\phi_h) - (\operatorname{div} \sigma_h,\phi_h) = (f,\phi_h), \quad \forall \ \phi_h \in \Lambda_h^n, \quad t \in I,$$

$$(u_h,\operatorname{div} \omega_h) + (\sigma_h,\omega_h) = 0, \quad \forall \ \omega_h \in \Lambda_h^{n-1}, \quad t \in I,$$

$$u_h(0) = g_h.$$
(4.2)

Define g_h to be the solution to the elliptic problem with load data $-\Delta g$, i.e.

$$\begin{aligned} (\operatorname{div} \hat{\sigma}_h, \phi_h) - (\Delta g, \phi_h) &= 0 \quad \forall \ \phi_h \in \Lambda_h^n, \\ (\hat{\sigma}_h, \omega_h) + (g_h, \operatorname{div} \omega_h) &= 0, \quad \forall \ \omega_h \in \Lambda_h^{n-1}. \end{aligned}$$

$$(4.3)$$

It is shown in [33] that a unique solution to (4.2) exists, based on the positive-definiteness of the solution operator $T_h : L^2 \to \Lambda_h^n$ for the elliptic problem. A more basic argument for this result can also be made by appealing to the existence of an adjoint to the discrete divergence operator.

Elliptic projection, an idea dating back to Wheeler [35], can be carried out for any fixed time value as we now discuss. For any $t_0 \in I$, define the **time-ignorant discrete** elliptic problem: Find $(\tilde{u}_h, \tilde{\sigma}_h) \in \Lambda_h^n \times \Lambda_h^{n-1}$ such that

$$(\operatorname{div} \tilde{\sigma}_{h}, \phi_{h}) + (-\Delta u(t_{0}), \phi_{h}) = 0, \quad \forall \phi_{h} \in \Lambda_{h}^{n},$$

$$(\tilde{\sigma}_{h}, \omega_{h}) + (\tilde{u}_{h}, \operatorname{div} \omega_{h}) = 0, \quad \forall \omega_{h} \in \Lambda_{h}^{n-1},$$

$$\tilde{u}_{h}(0) = g_{h}.$$
(4.4)

Note that the u appearing in the first equation of (4.4) is the solution to the continuous problem (4.1). Thus, we can view $\tilde{\sigma}_h$ and \tilde{u}_h as functions of t with the understanding that they are defined for each t value by (4.4) alone; no continuity with respect to t is required, hence the moniker 'time-ignorant.'

For ease of notation, and in keeping with Thomée, define the error functions

$$\rho(t) := \tilde{u}_h(t) - u(t),$$

$$\theta(t) := u_h(t) - \tilde{u}_h(t),$$

$$\varepsilon(t) := \sigma_h(t) - \tilde{\sigma}_h(t).$$

We now prove a lemma which will aid in our subsequent analysis. The result appears as part of the proof of Thomée [33, Theorem 17.2] but we expand it here for clarity.

Lemma 4.1 (Thomée [33]). *The error functions satisfy the semi-discrete formulation:*

$$\begin{aligned} &(\theta_t, \phi_h) - (\operatorname{div} \varepsilon, \phi_h) &= -(\rho_t, \phi_h), \quad \forall \ \phi_h \in \Lambda_h^n, \quad t \in I, \\ &(\theta, \operatorname{div} \omega_h) + (\varepsilon, \omega_h) &= 0, \qquad \forall \ \omega_h \in \Lambda_h^{n-1}, \quad t \in I. \end{aligned}$$

$$(4.5)$$

Proof. The second equation is immediate from the second equations in (4.2) and (4.4). The first equation can be written out as

$$(u_{h,t},\phi_h) - (\operatorname{div} \sigma_h,\phi_h) + (\operatorname{div} \tilde{\sigma}_h,\phi_h) - (\tilde{u}_{h,t},\phi_h) = (u_t,\phi_h) - (\tilde{u}_{h,t},\phi_h)$$

which is reduced as follows:

$$(u_{h,t},\phi_h) - (\operatorname{div} \sigma_h,\phi_h) + (\operatorname{div} \tilde{\sigma}_h,\phi_h) = (u_t,\phi_h)$$
 cancel like terms

$$(u_{h,t},\phi_h) - (\operatorname{div} \sigma_h,\phi_h) = -(\Delta u,\phi_h) + (u_t,\phi_h) \qquad \text{by (4.4)}$$

$$(f, \phi_h) = -(\Delta u, \phi_h) + (u_t, \phi_h)$$
 by (4.2)

This says that the continuous problem $u_t - \Delta u = f$ should hold in a weak sense when tested against any of the functions in Λ_h^n . This is guaranteed to be true since we chose $\Lambda_h^n \subset \Lambda^n = L^2$. Thus, the error equations hold as stated.

The following theorem says that if Λ_h^n and Λ_h^{n-1} are chosen according to the FEEC framework, then error estimates akin to (2.4) and (2.5) can be obtained. Note that in the semidiscrete setting, $(\Delta u)_t(t) = \partial_t \Delta u(t)$ since the time and spatial derivatives commute, allowing the simplified notation $\Delta u_t(t)$ used here.

Theorem 4.2. Fix $\Omega \subset \mathbb{R}^n$ and fix I := [0,T]. Suppose (u, σ) is the solution to (4.1) such that the regularity estimate

$$||u(t)||_{H^{s+2}} + ||\sigma(t)||_{H^{s+1}} \le c \, ||\Delta u(t)||_{H^s} \tag{4.6}$$

holds for $0 \leq s \leq s_{\max}$ and $t \in I$.

Choose finite element spaces

$$\Lambda_{h}^{n-1} = \left\{ \begin{array}{c} \mathcal{P}_{r+1}\Lambda^{n-1}(\mathcal{T}) \\ or \\ \mathcal{P}_{r+1}^{-}\Lambda^{n-1}(\mathcal{T}) \end{array} \right\}, \quad \Lambda_{h}^{n} = \mathcal{P}_{r+1}^{-}\Lambda^{n}(\mathcal{T}) \quad (= \mathcal{P}_{r}\Lambda^{n}(\mathcal{T}))$$

Then for $0 \le s \le s_{\text{max}}$, g_h defined by (4.3), and (u_h, σ_h) the solution to (4.2), the following error estimates hold:

$$||u_{h} - u||_{L^{2}(I, L^{2}\Lambda^{n})} \leq \begin{cases} ch\left(||\Delta u||_{L^{2}(I, L^{2})} + \sqrt{T} ||\Delta u_{t}||_{L^{1}(I, L^{2})}\right) & \text{if } r = 0\\ ch^{2+s}\left(||\Delta u||_{L^{2}(I, H^{s})} + \sqrt{T} ||\Delta u_{t}||_{L^{1}(I, H^{s})}\right) & \text{for } r > 0,\\ if s \leq r - 1 \\ (4.7) \end{cases}$$

$$||\sigma_{h} - \sigma||_{L^{2}(I,L^{2}\Lambda^{n-1})} \leq \begin{cases} ch\left(||\Delta u||_{L^{2}(I,H^{s})} + \sqrt{T} ||\Delta u_{t}||_{L^{2}(I,L^{2})}\right) \\ if r = 0, s = 0, \Lambda_{h}^{n-1} = \mathcal{P}_{1}^{-}\Lambda^{n-1}(\mathcal{T}) \\ c\left(h^{1+s} ||\Delta u||_{L^{2}(I,H^{s})} + h\sqrt{T} ||\Delta u_{t}||_{L^{2}(I,L^{2})}\right) \\ if r = 0, s \leq 1, \Lambda_{h}^{n-1} = \mathcal{P}_{1}\Lambda^{n-1}(\mathcal{T}) \\ c\left(h^{1+s} ||\Delta u||_{L^{2}(I,H^{s})} + h^{(3/2)+s}\sqrt{T} ||\Delta u_{t}||_{L^{2}(I,H^{s})}\right) \\ for r > 0, if s \leq r-1 \end{cases}$$
(4.8)

$$||\operatorname{div}(\sigma_{h} - \sigma)||_{L^{2}(I, L^{2}\Lambda^{n})} \leq \begin{cases} c\left(h^{s} ||\Delta u||_{L^{2}(I, H^{s})} + h ||\Delta u_{t}||_{L^{2}(I, L^{2})}\right) \\ if r = 0, s \leq 1 \\ c\left(h^{s} ||\Delta u||_{L^{2}(I, H^{s})} + h^{2+s} ||\Delta u_{t}||_{L^{2}(I, H^{s})}\right) \\ for r > 0, if s \leq r - 1 \end{cases}$$

$$(4.9)$$

Remark 4.3. Previous literature on semi-discrete methods usually leaves regularity assumptions implied by the error estimates. For instance, if $||u(t)||_{H^3}$ appears on the right side, it is implicitly assumed that $u(t) \in H^3$ for all $t \in I$. We have stated the specific regularity assumption (4.6) to make clear what regularity must be assumed and to follow the presentation from Arnold, Falk, and Winther [3, p. 342]. The careful reader will notice that the left side of (4.6) does not include a du(t) term, since $u(t) \in \Lambda^n$ implies du(t) = 0, nor a $||d\sigma(t)||_{H^s}$ term, since this can be absorbed into the $||\sigma(t)||_{H^{s+1}}$ term. An additional, more subtle difference is that f on the right side of (4.6) has been replaced by Δu . While these two are equivalent in the elliptic case, $\Delta u(t)$ evolves based on the initial data g while f(t) is prescribed, meaning they are in general different in the parabolic setting.

Proof. Observe that (4.4) is exactly the k = n case of the Hodge-Laplacian problem analyzed by Arnold, Falk and Winther [3] and the hypotheses here match their hypotheses. We can thus use a triangle inequality argument for each estimate, e.g.

$$||u(t) - u_h(t)|| \le ||u(t) - \tilde{u}_h(t)|| + ||\tilde{u}_h(t) - u_h(t)|| = ||\rho(t)|| + ||\theta(t)||$$
(4.10)

The first term will be bounded using the estimates from [3] and the second by the techniques from Thomée [33]. The FEEC estimate [3, p. 342] gives immediately

$$\|\rho(t)\| \le \begin{cases} ch \|\Delta u(t)\| & \text{if } r = 0\\ ch^{2+s} \|\Delta u(t)\|_{H^s} & \text{if } s \le r-1, \text{ for } r > 0 \end{cases}$$
(4.11)

Bounding $||\theta(t)||_{L^2}$ is more subtle. Set $\phi_h := \theta$ and $\omega_h := \varepsilon$ in (4.5). Adding the equations yields

$$\frac{1}{2}\frac{d}{dt}||\theta||^{2} + ||\varepsilon||^{2} = -(\rho_{t},\theta), \ t \in I$$
(4.12)

We use a technique from Thomée [33, p. 8] to derive an estimate for $||\theta(t)||$. Since $||\theta||$ may not be differentiable when $\theta = 0$, introduce a constant $\delta > 0$ and observe that

$$(||\theta||^{2} + \delta^{2})^{1/2} \frac{d}{dt} (||\theta||^{2} + \delta^{2})^{1/2} = \frac{1}{2} \frac{d}{dt} (||\theta||^{2} + \delta^{2}) = \frac{1}{2} \frac{d}{dt} ||\theta||^{2} \le ||\rho_{t}|| ||\theta||,$$

the last step following by (4.12) and Cauchy-Schwarz. Since $||\theta|| \le (||\theta||^2 + \delta^2)^{1/2}$, we have that

$$\frac{d}{dt}(||\theta||^2 + \delta^2)^{1/2} \le ||\rho_t||.$$

Note that $\theta(0) = u_h(0) - \tilde{u}_h(0) = g_h - g_h = 0$. Thus

$$||\theta(t)|| = \lim_{\delta \to 0} \int_0^t \frac{d}{dt} (||\theta||^2 + \delta^2)^{1/2} \le \int_0^t ||\rho_t||.$$

Using the bounds on $||\rho(t)||$ from (4.11), we get

$$||\theta(t)|| \le \begin{cases} ch \int_0^t ||\Delta u_t(\ell)|| d\ell & \text{if } r = 0, \\ ch^{2+s} \int_0^t ||\Delta u_t(\ell)||_{H^s} d\ell & \text{for } r > 0, \text{if } s \le r - 1. \end{cases}$$
(4.13)

We can now assemble estimate (4.7) by collecting our results. We show the technique of the case r = 0 as the other case employs identical analysis.

$$\begin{aligned} ||u_{h} - u||_{L^{2}(I,L^{2}\Lambda^{n})} &= \left(\int_{0}^{T} ||u_{h}(t) - u(t)||^{2} dt\right)^{1/2} \\ &\leq \left(\int_{0}^{T} (||\rho(t)|| + ||\theta(t)||)^{2} dt\right)^{1/2} \\ &\leq ch \left(\int_{0}^{T} \left(||\Delta u(t)|| + \int_{0}^{t} ||\Delta u_{t}(\ell)|| d\ell\right)^{2} dt\right)^{1/2} \\ &\leq ch \left(\int_{0}^{T} 2 \left(||\Delta u(t)||^{2} + \left(\int_{0}^{t} ||\Delta u_{t}(\ell)|| d\ell\right)^{2}\right) dt\right)^{1/2}. \end{aligned}$$

Roll the 2 into the constant c and observe that the inner integral is maximal when t = T. Thus,

$$\begin{aligned} ||u_{h} - u||_{L^{2}(I,L^{2}\Lambda^{n})} &\leq ch \left(\int_{0}^{T} ||\Delta u(t)||^{2} + ||\Delta u_{t}||_{L^{1}(I,L^{2})}^{2} dt \right)^{1/2} \\ &= ch \left(||\Delta u||_{L^{2}(I,L^{2})}^{2} + T ||\Delta u_{t}||_{L^{1}(I,L^{2})}^{2} \right)^{1/2} \\ &\leq ch \left(||\Delta u||_{L^{2}(I,L^{2})}^{2} + T ||\Delta u_{t}||_{L^{1}(I,L^{2})}^{2} + 2 ||\Delta u||_{L^{2}(I,L^{2})} \sqrt{T} ||\Delta u_{t}||_{L^{1}(I,L^{2})} \right)^{1/2} \\ &= ch \left(||\Delta u||_{L^{2}(I,L^{2})} + \sqrt{T} ||\Delta u_{t}||_{L^{1}(I,L^{2})} \right). \end{aligned}$$

We now turn to (4.8), i.e. an error bound for the approximation of σ . We use the same technique of bounding $||\sigma(t) - \tilde{\sigma}_h(t)||$ by the corresponding FEEC estimate and $||\tilde{\sigma}_h(t) - \sigma_h(t)|| = ||\varepsilon(t)||$ by a modification of (4.5). First, observe that the FEEC estimate [3, p. 342] gives

$$||\sigma(t) - \tilde{\sigma}_h(t)|| \le ch^{1+s} ||\Delta u(t)||_{H^s}, \text{ if } \begin{cases} s \le r+1, \quad \Lambda_h^{n-1} = \mathcal{P}_{r+1}\Lambda^{n-1}(\mathcal{T}) \\ s \le r, \qquad \Lambda_h^{n-1} = \mathcal{P}_{r+1}^-\Lambda^{n-1}(\mathcal{T}) \end{cases}$$

$$(4.14)$$

To bound $||\epsilon(t)||$, differentiate the second equation of (4.5) with respect to t and set $\phi_h := 2\theta_t, \omega_h := 2\varepsilon$, yielding

$$\begin{aligned} &(\theta_t, 2\theta_t) - (\operatorname{div} \varepsilon, 2\theta_t) &= -(\rho_t, 2\theta_t), \\ &(\varepsilon_t, 2\varepsilon) + 2(\theta_t, \operatorname{div} \varepsilon) &= 0. \end{aligned}$$

Adding the equations and converting to norms, we have the bound

$$\frac{d}{dt} ||\varepsilon||^2 + 2 ||\theta_t||^2 = -2(\rho_t, \theta_t) \le ||\rho_t||^2 + ||\theta_t||^2,$$

by Cauchy-Schwarz and the AM-GM inequality. Note that since $\theta(0) = 0$, we have $\varepsilon(0) = 0$ by the second equation of (4.5). Thus

$$||\varepsilon(t)||^{2} = \int_{0}^{t} \frac{d}{ds} ||\varepsilon(s)||^{2} ds \leq \int_{0}^{t} \frac{d}{ds} ||\varepsilon(s)||^{2} + ||\theta_{t}||^{2} ds \leq \int_{0}^{t} ||\rho_{t}||^{2} ds.$$
(4.15)

As before, we use (4.11) to derive

$$||\varepsilon(t)|| \leq \begin{cases} ch\left(\int_{0}^{t} ||\Delta u_{t}(\ell)||d\ell\right)^{1/2} & \text{if } r = 0, \\ ch^{2+s}\left(\int_{0}^{t} ||\Delta u_{t}(\ell)||_{H^{s}}^{2} d\ell\right)^{1/2} & \text{for } r > 0, \text{ if } s \leq r - 1. \end{cases}$$
(4.16)

We assemble estimate (4.8) by collecting these results in the same fashion we did for (4.7). In the r = 0 cases, we have

$$\begin{aligned} ||\sigma_{h} - \sigma||_{L^{2}(I,L^{2}\Lambda^{n-1})} &= \left(\int_{0}^{T} ||\sigma_{h}(t) - \sigma(t)||^{2} dt\right)^{1/2} \\ &\leq \left(\int_{0}^{T} (||\sigma(t) - \tilde{\sigma}_{h}(t)|| + ||\varepsilon(t)||)^{2} dt\right)^{1/2} \\ &\leq c \left(\int_{0}^{T} \left(h^{1+s} ||\Delta u(t)||_{H^{s}} + h \left(\int_{0}^{t} ||\Delta u_{t}(\ell)|| d\ell\right)^{1/2}\right)^{2} dt\right)^{1/2} \\ &\leq ch \left(\int_{0}^{T} 2 \left(h^{2s} ||\Delta u(t)||_{H^{s}}^{2} + \int_{0}^{t} ||\Delta u_{t}(\ell)|| d\ell\right) dt\right)^{1/2}. \end{aligned}$$

Rolling the 2 into the constant c and again noting that the inner integral is maximal when t = T, we recover the first two estimates of (4.8):

$$\begin{aligned} ||\sigma_{h} - \sigma||_{L^{2}(I,L^{2}\Lambda^{n-1})} &\leq ch \left(\int_{0}^{T} h^{2s} \left| |\Delta u(t)| \right|_{H^{s}}^{2} + \left| |\Delta u_{t}| \right|_{L^{2}(I,L^{2})}^{2} dt \right)^{1/2} \\ &= ch \left(h^{2s} \left| |\Delta u| \right|_{L^{2}(I,H^{s})}^{2} + T \left| |\Delta u_{t}| \right|_{L^{2}(I,L^{2})}^{2} dt \right)^{1/2} \\ &\leq ch \left(h^{s} \left| |\Delta u| \right|_{L^{2}(I,H^{s})}^{2} + \sqrt{T} \left| |\Delta u_{t}| \right|_{L^{2}(I,L^{2})}^{2} \right). \end{aligned}$$

When r > 0, (4.14) requires $s \le r$ or $s \le r + 1$ to obtain optimal convergence rates on the first term of the right side while (4.16) requires $s \le r - 1$ to obtain optimal rates on the second term. Thus the hypothesis $s \le r - 1$ implies both (4.14) and (4.16). The last estimate of (4.8) then follows by identical analysis to the first two cases.

Finally, we turn to estimate (4.9) and follow the same technique. Since div is the exterior derivative $d : \Lambda^{n-1} \to \Lambda^n$, we have from FEEC [3, p. 342] that

$$||\operatorname{div}(\sigma(t) - \tilde{\sigma}_h(t))|| \le ch^s \, ||\Delta u(t)||_{H^s}, \text{ if } s \le r+1.$$

$$(4.17)$$

Taking the L^2 norm with respect to time over [0, T] yields

$$\||\operatorname{div}(\sigma - \tilde{\sigma}_h)\|_{L^2(I, L^2)} \le ch^s \||\Delta u\|_{L^2(I, H^s)}, \text{ if } s \le r+1.$$
(4.18)

To bound $||\operatorname{div} \varepsilon||$ set $w_h := \varepsilon$ in (4.5) and take the derivative with respect to t. This yields

$$\frac{d}{dt} ||\varepsilon||^2 + (\theta_t, \operatorname{div} \varepsilon) = 0$$

Note that div $\varepsilon \in \Lambda_h^n$ since the discrete spaces are chosen to satisfy the relationship $\operatorname{div} \Lambda_h^{n-1} \subset \Lambda_h^n$. Thus, we can set $\phi_h := \operatorname{div} \varepsilon$ in in (4.5) and substitute to get

$$\frac{d}{dt} ||\varepsilon||^2 + ||\operatorname{div} \varepsilon||^2 = (\rho_t, \operatorname{div} \varepsilon).$$
(4.19)

Integrating both sides over [0, T] and applying Young's inequality, we have

$$||\varepsilon(T)||^{2} - ||\varepsilon(0)||^{2} + \int_{0}^{T} ||\operatorname{div} \varepsilon||^{2} dt \leq \frac{1}{2} \int_{0}^{T} ||\rho_{t}||^{2} + ||\operatorname{div} \varepsilon||^{2} dt$$
(4.20)

Note that $\varepsilon(0) = \sigma_h(0) - \tilde{\sigma}_h(0) = 0$ since the elliptic projection is the identity at time t = 0. We thus have

$$\int_{0}^{T} ||\operatorname{div} \varepsilon||^{2} dt \leq \int_{0}^{T} ||\rho_{t}||^{2} dt.$$
(4.21)

and by taking square roots,

$$\|\operatorname{div} \varepsilon\|_{L^{2}(I,L^{2})} \le \|\rho_{t}\|_{L^{2}(I,L^{2})} \, dt.$$
(4.22)

Again, we use (4.11) to get

$$\|\operatorname{div} \varepsilon\|_{L^{2}(I,L^{2})} \leq \begin{cases} ch \|\Delta u_{t}\|_{L^{2}(I,L^{2})} & \text{if } r = 0\\ ch^{2+s} \|\Delta u_{t}\|_{L^{2}(I,H^{s})} & \text{if } s \leq r-1, \text{ for } r > 0 \end{cases}$$

$$(4.23)$$

The estimate (4.9) follows by combining this with (4.18).

We now analyze hyperbolic problems again using the Bochner space norms introduced at the end of Section 3. Again, let $\Omega \subset \mathbb{R}^n$ be a contractible domain. Define the **velocity-stress mixed weak formulation**: Given the data functions f, u_0, u_1 , find $(\mu, \sigma) : I \to H\Lambda^n \times H\Lambda^{n-1}$ such that

$$\begin{aligned} (\mu_t, \phi) - (\operatorname{div} \sigma, \phi) &= (f, \phi), &\forall \phi \in H\Lambda^n, \quad t \in I, \\ (\mu, \operatorname{div} \omega) + (\sigma_t, \omega) &= 0, \quad \forall \omega \in H\Lambda^{n-1}, \quad t \in I, \\ \mu(0) &= u_1, \\ \sigma(0) &= \nabla u_0, \end{aligned}$$

$$(5.1)$$

where $\mu = u_t$ as in (2.8). The **semi-discrete hyperbolic problem** is thus: Find (μ_h, σ_h) : $I \to \Lambda_h^n \times \Lambda_h^{n-1}$ such that

$$(\mu_{h,t},\phi_h) - (\operatorname{div} \sigma_h,\phi_h) = (f,\phi_h), \quad \forall \phi_h \in \Lambda_h^n, \quad t \in I,$$

$$(\mu_h,\operatorname{div} \omega_h) + (\sigma_{h,t},\omega_h) = 0, \quad \forall \omega_h \in \Lambda_h^{n-1}, \quad t \in I,$$

$$\mu_h(0) = u_{1,h},$$

$$\sigma(0) = (\nabla u_0)_h.$$
(5.2)

We now generalize the results of Geveci [15] and others into the language of FEEC. We first prove a very simple proposition explaining the approximation properties of the π_h^k operators in this context.

Proposition 5.1. Choose finite element spaces

$$\Lambda_h^{n-1} = \left\{ \begin{array}{c} \mathcal{P}_{r+1}\Lambda^{n-1}(\mathcal{T}) \\ or \\ \mathcal{P}_{r+1}^-\Lambda^{n-1}(\mathcal{T}) \end{array} \right\}, \quad \Lambda_h^n = \mathcal{P}_{r+1}^-\Lambda^n(\mathcal{T}) \quad (= \mathcal{P}_r\Lambda^n(\mathcal{T}))$$

The smoothed projection operators from (3.1) have the approximation properties

$$\begin{aligned} \left| \left| \pi_{h}^{n-1} \omega - \omega \right| \right|_{L^{2} \Lambda^{n-1}} &\leq c \ h^{s} \left| \left| \omega \right| \right|_{H^{s} \Lambda^{n-1}}, \\ for \ 0 &\leq s \leq r+2, \quad \text{if } \Lambda_{h}^{n-1} = \mathcal{P}_{r+1} \Lambda^{n-1}(\mathcal{T}), \text{ or } \\ for \ 0 &\leq s \leq r+1, \quad \text{if } \Lambda_{h}^{n-1} = \mathcal{P}_{r+1}^{-} \Lambda^{n-1}(\mathcal{T}), \end{aligned}$$

$$(5.3)$$

$$||\pi_h^n \phi - \phi||_{L^2 \Lambda^n} \le c \, h^s \, ||\phi||_{H^s \Lambda^n} \,, \ \text{for } 0 \le s \le r+1.$$
(5.4)

Proof. Estimate (5.3) follows directly from Theorem 3.1 (*i*.). Note that Theorem 3.1 (*i*.) is stated for the case $\mathcal{P}_r \Lambda^{n-1}$ while here we have $\mathcal{P}_{r+1} \Lambda^{n-1}$, thereby allowing for the higher bound on *s* in this case. Finally, since $\pi_h^n \mu - \mu = \partial_t (\pi_h^n u - u)$, Theorem 3.1 (*i*.) also implies (5.4).

Theorem 5.2. Fix $\Omega \subset \mathbb{R}^n$ and fix I := [0, T]. Choose finite element spaces

$$\Lambda_h^{n-1} = \left\{ \begin{array}{c} \mathcal{P}_{r+1}\Lambda^{n-1}(\mathcal{T}) \\ or \\ \mathcal{P}_{r+1}^-\Lambda^{n-1}(\mathcal{T}) \end{array} \right\}, \quad \Lambda_h^n = \mathcal{P}_{r+1}^-\Lambda^n(\mathcal{T}) \quad (= \mathcal{P}_r\Lambda^n(\mathcal{T})) \,.$$

Then for (μ_h, σ_h) the solution to (5.2), the following error estimate holds:

$$||\mu_h - \mu||_{L^2(I, L^2\Lambda^n)} + ||\sigma_h - \sigma||_{L^2(I, L^2\Lambda^{n-1})} \le c \left(\sqrt{T}E_1 + h^s \left(\sqrt{T}E_2 + E_3\right)\right),$$
(5.5)

where

$$\begin{split} E_1 &= ||u_1 - u_{1,h}||_{L^2} + \\ &||(\nabla u_0) - (\nabla u_0)_h||_{L^2} & (error \ due \ to \ discretization \ of \ initial \ data) \\ E_2 &= ||u_1||_{H^s} + ||\nabla u_0||_{H^s} & (regularity \ of \ initial \ data) \\ E_3 &= ||u_t||_{L^2(I,H^s)} + ||\sigma||_{L^2(I,H^s)} & (regularity \ of \ continuous \ solution \ to \ (5.1)) \end{split}$$

Remark 5.3. This theorem strengthens and generalizes the result by Geveci [15] for n = 2 where L^2 projection is used instead of the smoothed projection operators π_h^k . An article by Makridakis [24] extended Geveci's results to n = 3 in the context of linear elastodynamics, however both papers had to assume the existence of finite element spaces and projections to them with certain properties. Our result here makes clear

what these spaces and projections should be in the unified language of FEEC. Moreover, the fact that the π_h^k operators are not the L^2 projection and hence not self-adjoint requires a revised proof technique that ultimately allows the removal of the error term $||u_{tt}||_{L^2(I,H^s)} + ||\sigma_t||_{L^2(I,H^s)}$ appearing in prior error bounds. Recent work by Falk and Winther [12, 13] presents local L^2 -stable cochain projectors that could potentially be used in place of the π_h^k operators.

Proof. Define $\Psi := \Lambda^n \times \Lambda^{n-1}$ with finite dimensional subspace $\Psi_h := \Lambda_h^n \times \Lambda_h^{n-1}$. Denote the components of an element $\psi_i \in \Psi$ by $\{\phi_i, \omega_i\}$. The L^2 inner product and norm on Ψ are

$$(\psi_1,\psi_2)_{\Psi} := (\phi_1,\phi_2)_{L^2} + (\omega_1,\omega_2)_{L^2} \text{ and } ||\psi||_{\Psi} := \sqrt{(\psi,\psi)_{\Psi}}.$$

Define a skew-symmetric bilinear form $a: \Psi \times \Psi \to \mathbb{R}$ by

$$a(\psi_1,\psi_2) := -(\operatorname{div} \omega_1,\phi_2)_{L^2} + (\phi_1,\operatorname{div} \omega_2)_{L^2}.$$

Let $\xi := (\mu, \sigma) \in \Psi$ be the solution to (5.1) and let $\psi := (\phi, \omega) \in \Psi$ be arbitrary. Then adding the equations of (5.1) yields

$$(\xi_t, \psi)_{\Psi} + a(\xi, \psi) = (f, \phi)_{L^2} \qquad \forall \, \psi \in \Psi.$$
(5.6)

Similarly, from (5.2) we get

$$(\xi_{h,t},\psi_h)_{\Psi} + a(\xi_h,\psi_h) = (f,\phi_h)_{L^2} \qquad \forall \ \psi_h \in \Psi_h.$$
(5.7)

Define a projection operator $\pi_h : \Psi \to \Psi_h$ using the bounded cochain projections from (3.1) via $\pi_h \psi := \{\pi_h^n \phi, \pi_h^{n-1} \omega\}$. Since π_h only affects the spatial variables, it commutes with the time derivative operator, i.e. $(\partial_t \pi_h \xi, \psi_h)_{\Psi} = (\pi_h \partial_t \xi, \psi_h)_{\Psi}$. Using this and (5.6), and letting \mathbb{I} denote the identity operator, we derive

$$a(\pi_{h}\xi,\psi_{h}) = a(\xi,\psi_{h}) + a((\pi_{h} - \mathbb{I})\xi,\psi_{h})$$

$$(\partial_{t}\pi_{h}\xi,\psi_{h})_{\Psi} + a(\pi_{h}\xi,\psi_{h}) = a(\xi,\psi_{h}) + (\pi_{h}\partial_{t}\xi,\psi_{h})_{\Psi} + a((\pi_{h} - \mathbb{I})\xi,\psi_{h})$$

$$(\partial_{t}\pi_{h}\xi,\psi_{h})_{\Psi} + a(\pi_{h}\xi,\psi_{h}) = (f,\phi_{h})_{L^{2}} + ((\pi_{h} - \mathbb{I})\partial_{t}\xi,\psi_{h})_{\Psi} + a((\pi_{h} - \mathbb{I})\xi,\psi_{h}),$$

(5.8)

which holds for all $\psi_h \in \Psi_h \subset \Psi$. Now define the error function

$$\varepsilon_h(t) := \pi_h \xi(t) - \xi_h(t).$$

The derivation of a good bound for $||\varepsilon_h(t)||_{\Psi}$ constitutes the bulk of the remainder of the proof. Subtracting (5.7) from (5.8) yields

$$(\partial_t \varepsilon_h(t), \psi_h)_{\Psi} + a(\varepsilon_h(t), \psi_h) = ((\pi_h - \mathbb{I})\partial_t \xi(t), \psi_h)_{\Psi} + a((\pi_h - \mathbb{I})\xi(t), \psi_h), \quad \forall \, \psi_h \in \Psi_h.$$
(5.9)

Define the skew-adjoint linear operator $L_h: \Psi_h \to \Psi_h$ by

$$(L_h\psi_1,\psi_2)_{\Psi} := a(\psi_1,\psi_2) \qquad \forall \ \psi_1,\psi_2 \in \Psi_h.$$

We can thus re-write (5.9) as an equation of functionals on Ψ_h :

$$\partial_t \varepsilon_h(t) + L_h \varepsilon_h(t) = (\pi_h - \mathbb{I}) \partial_t \xi(t) + L_h(\pi_h - \mathbb{I}) \xi(t)$$
(5.10)

To ease notation, set $Q(t) := (\pi_h - \mathbb{I})\partial_t \xi(t)$ and $R(t) := (\pi_h - \mathbb{I})\xi(t)$, yielding

$$\partial_t \varepsilon_h(t) + L_h \varepsilon_h(t) = Q(t) + L_h R(t)$$
(5.11)

We will use some basic results from the theory of semigroups of linear operators as can be found, for instance, in [27]. For any fixed $\tau \in \mathbb{R}$, the product rule in this context yields

$$\partial_t (e^{(t-\tau)L_h}(\varepsilon_h(t) - R(t))) = L_h e^{(t-\tau)L_h}(\varepsilon_h(t) - R(t)) + e^{(t-\tau)L_h}\partial_t(\varepsilon_h(t) - R(t))$$
$$= e^{(t-\tau)L_h} (\partial_t \varepsilon_h(t) + L_h \varepsilon_h(t) - (L_h + \partial_t)R(t))$$

Note that we used the fact that $e^{(t-\tau)L_h}$ commutes with L_h , a standard result [27, Corollary 1.4]. Swapping the roles of t and τ , we re-write the above as

$$\partial_{\tau}(e^{-(t-\tau)L_h}(\varepsilon_h(\tau) - R(\tau))) = e^{-(t-\tau)L_h}\left(\partial_{\tau}\varepsilon_h(\tau) + L_h\varepsilon_h(\tau) - (L_h + \partial_{\tau})R(\tau)\right)$$
(5.12)

Now we integrate in such a way that (5.11) and (5.12) will give us an expression for $\varepsilon_h(t)$. First observe that $\partial_\tau R(\tau) = Q(\tau)$ since ∂_τ commutes with π_h and \mathbb{I} . Thus,

$$0 = \int_0^t e^{-(t-\tau)L_h} \left(Q(\tau) - \partial_\tau R(\tau) \right) d\tau$$

=
$$\int_0^t e^{-(t-\tau)L_h} \left(Q(\tau) + L_h R(\tau) - (L_h + \partial_\tau) R(\tau) \right) d\tau \qquad \text{by } \pm L_h R(\tau)$$

$$= \int_0^\tau e^{-(t-\tau)L_h} \left(\partial_\tau \varepsilon_h(\tau) + L_h \varepsilon_h(\tau) - (L_h + \partial_\tau) R(\tau) \right) d\tau \qquad \text{by (5.11)}$$

$$= \int_0^t \partial_\tau (e^{-(t-\tau)L_h}(\varepsilon_h(\tau) - R(\tau)) d\tau \qquad \qquad \text{by (5.12)}$$

$$=\varepsilon_h(t) - R(t) - e^{-tL_h}\varepsilon_h(0) + R(0)$$
 fund. thm. calculus

Rewriting the above chain of equalities, we see that

$$\varepsilon_h(t) = e^{-tL_h} \varepsilon_h(0) + R(t) - R(0)$$
(5.13)

Observe that e^{-tL_h} is unitary meaning it preserves Ψ -norm, i.e. $||e^{-tL_h}\psi||_{\Psi} = ||\psi||_{\Psi}$ for all $\psi \in \Psi$. This follows from the fact that L_h is a real, skew self-adjoint operator, meaning iL_h is self-adjoint, which is equivalent to saying e^{-tL_h} is unitary [27, Theorem 10.8]. Thus, taking the $||\cdot||_{\Psi}$ norm of (5.13), the triangle inequality gives

$$\begin{aligned} ||\varepsilon_{h}(t)||_{\Psi} &\leq ||\varepsilon_{h}(0)||_{\Psi} + ||R(t)||_{\Psi} + ||R(0)||_{\Psi} \\ &= ||\varepsilon_{h}(0)||_{\Psi} + ||(\pi_{h} - \mathbb{I})\xi(t)||_{\Psi} + ||(\pi_{h} - \mathbb{I})\xi(0)||_{\Psi} , \tag{5.14}$$

Unpacking the notation lets us characterize this bound in terms of the errors defined in the theorem statement. Recall that u_0 and u_1 are given initial data functions and should not be confused with u_h or u_t . We will use $f \leq g$ to mean $f \leq cg$ where c is some constant independent of h and T. In $||\cdot||_{\Psi}$ norm, we then have

$$||\varepsilon_{h}(t)|| \lesssim ||\pi_{h}^{n}u_{1} - u_{1,h}|| + \left| \left| \pi_{h}^{n-1}(\nabla u_{0}) - (\nabla u_{0})_{h} \right| \right| + ||(\pi_{h} - \mathbb{I})\xi(t)|| + ||(\pi_{h} - \mathbb{I})\xi(0)||$$
(5.15)

To bound the first term on the right, use (5.4) from Proposition 5.1 to get

$$||\pi_h^n u_1 - u_{1,h}|| \le ||\pi_h^n u_1 - u_1|| + ||u_1 - u_{1,h}|| \lesssim h^s ||u_1||_{H^s} + ||u_1 - u_{1,h}||$$

Using (5.3) likewise for the second term, we have

$$\left|\left|\pi_{h}^{n}u_{1}-u_{1,h}\right|\right|+\left|\left|\pi_{h}^{n-1}(\nabla u_{0})-(\nabla u_{0})_{h}\right|\right| \lesssim E_{1}+h^{s}E_{2}$$
(5.16)

Also by (5.3) and (5.4), we have the bounds

$$||(\pi_h - \mathbb{I})\xi(t)|| \lesssim h^s(||\sigma(t)||_{H^s\Lambda^{n-1}} + ||u_t(t)||_{H^s\Lambda^n})$$
(5.17)

$$\|(\pi_h - \mathbb{I})\xi(0)\| \lesssim h^s(\|\nabla u_0\|_{H^s\Lambda^{n-1}} + \|u_1\|_{H^s\Lambda^n})$$
(5.18)

Using (5.15) in conjunction with (5.16), (5.17), and (5.18), we derive

$$\int_{0}^{T} ||\varepsilon_{h}(t)||^{2} dt \lesssim TE_{1}^{2} + h^{2s}TE_{2}^{2} + h^{2s} \int_{0}^{T} ||\sigma(t)||_{H^{s}}^{2} + ||u_{t}(t)||_{H^{s}}^{2} + E_{2}^{2} dt$$
$$\lesssim TE_{1}^{2} + h^{2s}TE_{2}^{2} + h^{2s}E_{3} + h^{2s}TE_{2}^{2}$$
(5.19)

We now start building up the main result.

$$\left(||\mu_{h} - \mu||_{L^{2}(I,L^{2}\Lambda^{n})} + ||\sigma_{h} - \sigma||_{L^{2}(I,L^{2}\Lambda^{n-1})} \right)^{2}$$

$$\lesssim \int_{0}^{T} \left(||\mu_{h} - \pi_{h}^{n}\mu|| + ||\sigma_{h} - \pi_{h}^{n-1}\sigma|| \right)^{2} + ||\pi_{h}^{n}\mu - \mu||^{2} + ||\pi_{h}^{n-1}\sigma - \sigma||^{2} dt$$

$$= \int_{0}^{T} ||\varepsilon_{h}(t)||^{2} dt + ||\pi_{h}^{n}\mu - \mu||^{2}_{L^{2}(I,L^{2}\Lambda^{n})} + ||\pi_{h}^{n-1}\sigma - \sigma||^{2}_{L^{2}(I,L^{2}\Lambda^{n-1})} dt$$

$$\lesssim \int_{0}^{T} ||\varepsilon_{h}(t)||^{2} + h^{2s} ||\mu(t)||^{2}_{H^{s}} + h^{2s} ||\sigma(t)||^{2}_{H^{s}} dt$$

$$\lesssim \int_{0}^{T} ||\varepsilon_{h}(t)||^{2} dt + h^{2s} E_{3}^{2}.$$

$$(5.20)$$

Combining (5.20) and (5.19) yields

$$\left(||\mu_h - \mu||_{L^2(I, L^2\Lambda^n)} + ||\sigma_h - \sigma||_{L^2(I, L^2\Lambda^{n-1})} \right)^2 \lesssim \left(TE_1^2 + 2h^{2s}TE_2^2 + 2h^{2s}E_3^2 \right)$$

$$\lesssim \left(\sqrt{T}E_1 + h^s \left(\sqrt{T}E_2 + E_3 \right) \right)^2$$

Taking the square root of both sides completes the proof.

6. SEMI-LINEAR EVOLUTION PROBLEMS

We now show how the techniques developed above can be extended to certain types of non-linear evolution problems. Consider the **semi-linear heat equation**: Find u(x, t) such that

$$u_t - \Delta u + F(u) = f \quad \text{in } \Omega, \quad \text{for } t > 0$$

$$u = 0 \quad \text{on } \partial\Omega, \quad \text{for } t > 0 \quad \text{with } u(\cdot, 0) = g \text{ in } \Omega, \tag{6.1}$$

where F is some non-linear operator on $L^2(\Omega)$. The existence and uniqueness of solutions to instances of this problem have been studied extensively [16, 18, 30, 36] as have finite element methods for the approximation of its solution [9, 23, 31, 33, 34].

We define the **semi-linear mixed weak form parabolic** problem: Given f and g, find $(u, \sigma) : I \to H\Lambda^n \times H\Lambda^{n-1}$ such that

$$(u_t, \phi) - (\operatorname{div} \sigma, \phi) + (F(u), \phi) = (f, \phi), \quad \forall \phi \in H\Lambda^n, \quad t \in I,$$

$$(u, \operatorname{div} \omega) + (\sigma, \omega) = 0, \quad \forall \omega \in H\Lambda^{n-1}, \quad t \in I,$$

$$u(0) = g.$$
(6.2)

The semi-linear semi-discrete parabolic problem is thus: Find $(u_h, \sigma_h) : I \to \Lambda_h^n \times \Lambda_h^{n-1}$ such that

$$(u_{h,t},\phi_h) - (\operatorname{div} \sigma_h,\phi_h) + (F(u_h),\phi_h) = (f,\phi_h), \quad \forall \phi_h \in \Lambda_h^n, \quad t \in I,$$

$$(u_h,\operatorname{div} \omega_h) + (\sigma_h,\omega_h) = 0, \quad \forall \omega_h \in \Lambda_h^{n-1}, \quad t \in I, \quad (6.3)$$

$$u_h(0) = g_h,$$

where $g_h \in \Lambda_h^n$ is an approximation of g. Analogously to the linear case, for any $t_0 \in I$, define the **time-ignorant linear discrete elliptic problem**: find $(\tilde{u}_h, \tilde{\sigma}_h) \in \Lambda_h^n \times \Lambda_h^{n-1}$ such that

$$(\operatorname{div} \tilde{\sigma}_{h}, \phi_{h}) - (\Delta u(t_{0}), \phi_{h}) = 0, \quad \forall \phi_{h} \in \Lambda_{h}^{n}, (\tilde{\sigma}_{h}, \omega_{h}) + (\tilde{u}_{h}, \operatorname{div} \omega_{h}) = 0, \quad \forall \omega_{h} \in \Lambda_{h}^{n-1}, \tilde{u}_{h}(0) = g_{h},$$

$$(6.4)$$

where now u is the solution to the continuous semi-linear problem (6.2). Recall, as in (4.4), that no continuity of $\tilde{\sigma}_h$ or \tilde{u}_h with respect to t is required. Similarly, define

$$\rho(t) := \tilde{u}_h(t) - u(t),$$

$$\theta(t) := u_h(t) - \tilde{u}_h(t),$$

$$\varepsilon(t) := \sigma_h(t) - \tilde{\sigma}_h(t),$$

where u, σ and their discrete counterparts are now solutions to the corresponding semilinear problems. We also introduce the error term

$$\eta(t) := F(u(t)) - F(u_h(t)).$$

Here, we assume that the nonlinear function $F : \mathbb{R} \to \mathbb{R}$ satisfies the following (local) Lipschitz condition:

$$(F(u) - F(u_h), v) \le K ||u - u_h||_{L^2} ||v||, \quad \forall v \in L^2(\Omega),$$
(6.5)

where u and u_h are the solutions to (6.2) and (6.3) respectively. We remark that this type of local Lipschitz condition is slightly different from those assumed by Holst and Stern in [21, 5]. Here, we use the L^2 norm on the right sides of the inequalities. By the Poincaré inequality, (6.5) implies the Lipschitz assumption used in [21, 5]. The key here is that we only require the assumption (6.5) to hold for the solutions u and u_h . For example, suppose that the solutions u and u_h satisfy some a priori L^{∞} bounds i.e. there exist some constants α and β such that $\alpha \leq u$, $u_h \leq \beta$. Then the local Lipschitz condition is satisfied if there is some constant K > 0 such that $|F'(\xi)| \leq K$ for all $\xi \in [\alpha, \beta]$.

We have the follow lemma, which reduces to Lemma 4.1 in the linear case.

Lemma 6.1. The semi-linear error functions satisfy the semi-discrete formulation, i.e.

$$\begin{aligned} & (\theta_t, \phi_h) - (\operatorname{div} \varepsilon, \phi_h) &= (\eta, \phi_h) - (\rho_t, \phi_h), \quad \forall \ \phi_h \in \Lambda_h^n, \quad t \in I, \\ & (\theta, \operatorname{div} \omega_h) + (\varepsilon, \omega_h) &= 0, \qquad \qquad \forall \ \omega_h \in \Lambda_h^{n-1}, \quad t \in I. \end{aligned}$$

$$(6.6)$$

Proof. The second equation is immediate from the second equations in (6.3) and (6.4). The first equation is derived as follows.

$$(u_t, \phi_h) - (\Delta u, \phi_h) + (F(u), \phi_h) = (f, \phi_h)$$
 by (6.1),

$$(u_t, \phi_h) - (\Delta u, \phi_h) + (F(u), \phi_h) = (u_{h,t}, \phi_h) - (\operatorname{div} \sigma_h, \phi_h) + (F(u_h), \phi_h)$$
 by (6.3)

$$(u_t, \phi_h) - (\operatorname{div} \tilde{\sigma}_h, \phi_h) + (F(u), \phi_h) = (u_{h,t}, \phi_h) - (\operatorname{div} \sigma_h, \phi_h) + (F(u_h), \phi_h)$$
 by (6.4),

from which the result follows by adding $(\pm \tilde{u}_{h,t}, \phi_h)$ and collecting terms.

Theorem 6.2. Fix $\Omega \subset \mathbb{R}^n$ and fix I := [0,T]. Suppose (u,σ) is the solution to (6.2) such that the regularity estimate

$$||u(t)||_{H^{s+2}} + ||\sigma(t)||_{H^{s+1}} + ||d\sigma(t)||_{H^s} \le c ||\Delta u(t)||_{H^s}$$
(6.7)

holds for $0 \le s \le s_{\text{max}}$ and $t \in I$. Assume that the operator F satisfies the Lipschitz assumption (6.5). Choose finite element spaces

$$\Lambda_h^{n-1} = \left\{ \begin{array}{c} \mathcal{P}_{r+1}\Lambda^{n-1}(\mathcal{T}) \\ or \\ \mathcal{P}_{r+1}^-\Lambda^{n-1}(\mathcal{T}) \end{array} \right\}, \quad \Lambda_h^n = \mathcal{P}_{r+1}^-\Lambda^n(\mathcal{T}) \quad (= \mathcal{P}_r\Lambda^n(\mathcal{T}))$$

Then for $0 \le s \le s_{\text{max}}$, g_h defined by (4.3), and (u_h, σ_h) the solution to (6.3), the following error estimates hold:

$$||u_{h} - u||_{L^{2}(I,L^{2}\Lambda^{n})} \lesssim \begin{cases} h\left(||\Delta u||_{L^{2}(I,L^{2})} + ||\Delta u_{t}||_{L^{2}(I,L^{2})}\right) & \text{if } r = 0\\ h^{1+s}\left(||\Delta u||_{L^{2}(I,H^{s})} + ||\Delta u_{t}||_{L^{2}(I,H^{s})}\right) & \text{for } r > 0, \\ \text{if } s \leq r - 1 \end{cases}$$

$$||\sigma_{h} - \sigma||_{L^{2}(I,L^{2}\Lambda^{n-1})} \lesssim \begin{cases} h\left(||\Delta u||_{L^{2}(I,H^{s})} + ||\Delta u_{t}||_{L^{2}(I,L^{2})}\right) \\ \text{if } r = 0, \ s = 0, \ \Lambda_{h}^{n-1} = \mathcal{P}_{1}^{-}\Lambda^{n-1}(\mathcal{T}) \\ h^{1+s} ||\Delta u||_{L^{2}(I,H^{s})} + h ||\Delta u_{t}||_{L^{2}(I,L^{2})} \\ \text{if } r = 0, \ s \leq 1, \ \Lambda_{h}^{n-1} = \mathcal{P}_{1}\Lambda^{n-1}(\mathcal{T}) \\ h^{1+s} ||\Delta u||_{L^{2}(I,H^{s})} + h^{(3/2)+s} ||\Delta u_{t}||_{L^{2}(I,H^{s})} \\ \text{for } r > 0 \ \text{if } s \leq r - 1 \end{cases}$$

$$(6.9)$$

$$\begin{cases} for r > 0, if s \le r - 1 \\ \|\Delta u\|_{L^{2}(I, L^{2}\Lambda^{n})} \lesssim \begin{cases} h^{s} \|\Delta u\|_{L^{2}(I, H^{s})} + h\| \Delta u_{t}\|_{L^{2}(I, L^{2})} \\ if r = 0, s \le 1 \\ h^{s} \|\Delta u\|_{L^{2}(I, H^{s})} + h^{2+s} \|\Delta u_{t}\|_{L^{2}(I, H^{s})} \\ for r > 0, if s \le r - 1 \end{cases}$$

$$(6.10)$$

Proof. The proof is very similar to that of Theorem 4.2. Equation (6.4) is the k = n case of the discrete mixed variational problem examined by Holst and Stern in [21, Equation (9)]. Therefore, we can use the same type of triangle inequality from (4.10) to recover the estimates. By [21, Theorem 4.2], we have the estimates

$$\|\rho(t)\| \le \begin{cases} ch \|\Delta u(t)\| & \text{if } r = 0\\ ch^{2+s} \|\Delta u(t)\|_{H^s} & \text{if } s \le r-1, \text{ for } r > 0 \end{cases}$$
(6.11)

$$\|\rho_t(t)\| \le \begin{cases} ch \|\Delta u_t(t)\| & \text{if } r = 0\\ ch^{2+s} \|\Delta u_t(t)\|_{H^s} & \text{if } s \le r-1, \text{ for } r > 0 \end{cases}$$
(6.12)

$$\|\sigma(t) - \tilde{\sigma}_h(t)\| \le ch^{1+s} \|\Delta u(t)\|_{H^s}, \text{ if } \begin{cases} s \le r+1, \quad \Lambda_h^{n-1} = \mathcal{P}_{r+1}\Lambda^{n-1}(\mathcal{T}) \\ s \le r, \qquad \Lambda_h^{n-1} = \mathcal{P}_{r+1}^-\Lambda^{n-1}(\mathcal{T}) \\ (6.13) \end{cases}$$

$$\|\operatorname{div}(\sigma(t) - \tilde{\sigma}_h(t))\| \le ch^s \, \|\Delta u(t)\|_{H^s}, \text{ if } s \le r+1.$$
(6.14)

An explanation of how these estimates are derived from the results of [21] is given in Appendix A. Note that these estimates are exactly the same as the corresponding estimates (4.11), (4.14) and (4.18) from the linear case.

To bound $||\theta||$, we adapt the proof of Theorem 4.2 to account for the non-linearity. Set $\phi_h := \theta$ and $\omega_h := \varepsilon$ in (6.6). Adding the equations yields

$$\frac{1}{2}\frac{d}{dt}\|\theta\|^2 + \|\varepsilon\|^2 + (F(u_h) - F(u), \theta) = -(\rho_t, \theta), \ t \in I$$
(6.15)

By the Lipschitz condition (6.5), we obtain:

$$\frac{1}{2} \frac{d}{dt} \|\theta\|^{2} \leq |(F(u) - F(u_{h}), \theta)| + |(\rho_{t}, \theta)|$$

$$\leq K \|u - u_{h}\| \|\theta\| + \|\rho_{t}\| \|\theta\|$$

$$\leq K \|\rho\| \|\theta\| + \|\rho_{t}\| \|\theta\| + K \|\theta\|^{2}$$

$$\leq C \|\theta\|^{2} + C(\|\rho\|^{2} + \|\rho_{t}\|^{2}).$$

Notice that $\theta(0) = 0$ and an application of Gronwall's inequality implies that

$$\begin{aligned} \|\theta(t)\|^2 &\leq 2Ce^{2Ct} \int_0^t e^{-2Cs} (\|\rho\|^2 + \|\rho_s\|^2) ds \\ &\leq 2Ce^{2Ct} \left(\|\rho\|_{L^2(I,L^2)}^2 + \|\rho_t\|_{L^2(I,L^2)}^2 \right) \end{aligned}$$

To get an estimate on $\|\theta\|_{L^2(I,L^2)}$, we integrate by t on both sides to obtain

$$\|\theta\|_{L^2(I,L^2)} \lesssim \sqrt{T}(\|\rho\|_{L^2(I,L^2)} + \|\rho_t\|_{L^2(I,L^2)}).$$
(6.16)

The estimate (6.8) then follows by (6.11), (6.12), (6.16) and a triangle inequality.

To obtain the error bound (6.9) for the approximation of σ , we use the same technique as in Theorem 4.2. To bound $||\epsilon(t)||$, we differentiate the second equation of (6.6) with respect to t and set $\phi_h := 2\theta_t$, $\omega_h := 2\varepsilon$, yielding

$$\begin{aligned} &(\theta_t, 2\theta_t) - (\operatorname{div} \varepsilon, 2\theta_t) &= (\eta, 2\theta_t) - (\rho_t, 2\theta_t), \\ &(\varepsilon_t, 2\varepsilon) + (\theta_t, 2\operatorname{div} \varepsilon) &= 0. \end{aligned}$$

Adding the equations and using the Lipschitz assumption (6.5), we have the bound

$$\frac{d}{dt} ||\varepsilon||^2 + 2 ||\theta_t||^2 = 2(\eta, \theta_t) - 2(\rho_t, \theta_t) \\ \leq 2K ||u - u_h|| ||\theta_t|| + 2 ||\rho_t|| ||\theta_t||$$

Now write $||u - u_h|| = ||\theta + \rho||$, apply the triangle inequality, and then apply Young's inequality to get a constant C such that

$$\frac{d}{dt} ||\varepsilon||^2 + 2 ||\theta_t||^2 \le C \left(||\rho||^2 + ||\theta||^2 + ||\rho_t||^2 \right) + ||\theta_t||^2.$$

Canceling one of the $\|\theta_t\|^2$ on both sides, and by the estimates (6.11)-(6.12) and (6.16), we obtain

$$\begin{aligned} ||\varepsilon(t)||^{2} &= \int_{0}^{t} \frac{d}{ds} ||\varepsilon(s)||^{2} ds \leq \int_{0}^{t} \frac{d}{ds} ||\varepsilon(s)||^{2} + ||\theta_{t}||^{2} ds \\ &\leq C \int_{0}^{t} ||\rho||^{2} + ||\theta||^{2} + ||\rho_{t}||^{2} ds. \end{aligned}$$

Now use the estimates (6.11) and (6.12) to derive

$$||\varepsilon(t)|| \leq \begin{cases} ch\sqrt{T} \left(\int_0^t ||\Delta u(\ell)|| + ||\Delta u_t(\ell)|| d\ell \right)^{1/2} & \text{if } r = 0, \\ ch^{2+s}\sqrt{T} \left(\int_0^t ||\Delta u(\ell)||_{H^s}^2 + ||\Delta u_t(\ell)||_{H^s}^2 d\ell \right)^{1/2} & \text{for } r > 0, \text{ if } s \leq r - 1 \end{cases}$$
(6.17)

The estimate (6.9) then follows from triangle inequality, (6.13), and (6.17).

Finally, to estimate (6.10), we follow the same techniques. We first bound $||\text{div }\varepsilon||$ by setting $w_h := \varepsilon$ in (6.6) and take the derivative with respect to t. This yields

$$\frac{d}{dt} \left| \left| \varepsilon \right| \right|^2 + (\theta_t, \operatorname{div} \varepsilon) = 0.$$

Note that div $\varepsilon \in \Lambda_h^n$ since the discrete spaces are chosen to satisfy the relationship $\operatorname{div} \Lambda_h^{n-1} \subset \Lambda_h^n$. Thus, we can set $\phi_h := \operatorname{div} \varepsilon$ in (6.6) and substitute to get

$$\frac{d}{dt} ||\varepsilon||^2 + ||\operatorname{div} \varepsilon||^2 = (\rho_t, \operatorname{div} \varepsilon) - (\eta, \operatorname{div} \varepsilon).$$
(6.18)

Integrating both sides over [0, T] and applying the Lipschitz assumption (6.5) and Young's inequality, we have

$$||\varepsilon(T)||^{2} - ||\varepsilon(0)||^{2} + \int_{0}^{T} ||\operatorname{div} \varepsilon||^{2} dt \leq C \int_{0}^{T} ||\rho_{t}||^{2} + \|\rho\|^{2} + \|\theta\|^{2} dt + \frac{1}{2} \int_{0}^{T} ||\operatorname{div} \varepsilon||^{2} dt.$$
(6.19)

Note that $\varepsilon(0) = \sigma_h(0) - \tilde{\sigma}_h(0) = 0$ since the elliptic projection is the identity at time t = 0. We thus have

$$\int_0^T ||\operatorname{div} \varepsilon||^2 dt \le 2C \int_0^T ||\rho_t||^2 + \|\rho\|^2 + \|\theta\|^2 dt.$$
(6.20) of follows similarly.

The rest of the proof follows similarly.

We remark that the estimates for the semi-linear case given by Theorem 6.2 are very similar to those for the linear case given by Theorem 4.2, although the latter are somewhat sharper. In the semi-linear case, the constant on the right side of the equation will have a factor of \sqrt{T} , while in the linear case the dependence on \sqrt{T} is made explicit. Further note that the estimate on $||u_h - u||$ in the linear case involves an $L^1(\cdot \cdot)$ norm on Δu_t as part of the bound, which is a stronger result than the bound we obtained in the semi-linear case that involves the $L^2(\cdot \cdot)$ norm on Δu_t .

7. CONCLUDING REMARKS

In this article, we have extended the Finite Element Exterior Calculus of Arnold, Falk, and Winther [3, 2] for linear mixed variational problems to linear and semi-linear parabolic and hyperbolic evolution systems. Both the parabolic and hyperbolic cases make strong use of the smoothed projection operators π_h^k , which are one of the most elaborate and delicate constructions in the FEEC framework. In the parabolic case, the use of the π_h^k operators was hidden somewhat by the use of elliptic projection error estimates, proofs of which rely on properties of these operators. In the hyperbolic case, the proof techniques use these properties more explicitly. In both cases, the formal treatment and generalization of these operators by Arnold, Falk and Winther can now be seen as a useful tool for the analysis of evolution problems as well as elliptic PDE. We have also seen in this article how the recent generalizations of the FEEC by Holst and Stern [20, 21] for semi-linear elliptic PDE can be extended to evolution PDE as well, both parabolic and hyperbolic types. We also anticipate that the basic approach to analyzing variational crimes in [20, 21] for the linear and semilinar elliptic cases will also work in the case of evolution problems; we will explore the question of variational crimes in a subsequent article, with the target being the analysis of surface finite element methods for evolution problems.

APPENDIX A. EXPLANATION OF SEMI-LINEAR ERROR ESTIMATES

In this appendix, we explain why estimates (6.11), (6.13), and (6.14) follow from [21, Theorem 4.2]. We will focus just on the r > 0 case of (6.11) as it requires the sharpening of a special case of an estimate appearing in [21, Theorem 4.2]. The other cases work out along similar lines by a direct application of the Holst and Stern estimates.

First, we recall some notation from [3] used in [21]. If (W, d) is a Hilbert complex with associated domain complex (V, d) and parametrized subcomplex family (V_h, d) , denote the best approximation in W-norm by

$$E(w) = \inf_{v \in V_h^k} ||w - v||_W, \quad w \in W^k.$$

The relevant result from Holst and Stern [21, Theorem 4.2] is stated as

$$||u - \tilde{u}_h||_V + ||p - p_h||_W \le c(E(u) + E(du) + E(p) + \eta[E(\sigma) + E(d\sigma)] + (\delta + \mu)E(d\sigma) + \mu E(P_{\mathfrak{B}}u)),$$
(A.1)

where η , δ , and μ are coefficients defined as the norms of certain abstract operators, $u \in W_k$, and p is a harmonic k-form with discrete counterpart p_h introduced to make the abstract Hodge-Laplacian problem well-posed.

Casting this into the context of the deRham complex, we have

 $(W, d) = (L^2\Lambda, d)$ and $(V, d) = (H\Lambda, d)$.

Since we are interested here only in the case k = n, there are no harmonic k-forms so that $p = p_h = 0$. Further, du = 0 since $d\Lambda^n = 0$, whereby $||u - \tilde{u}_h||_V = ||u - \tilde{u}_h||_W = ||u - \tilde{u}_h||_{L^2}$. This eliminates the error terms in p and du, giving us the reduced estimate

$$||u - \tilde{u}_h||_{L^2} \le c(E(u) + \eta[E(\sigma) + E(d\sigma)] + (\delta + \mu)E(d\sigma) + \mu E(P_{\mathfrak{B}}u)).$$

Crucially, this estimate can be reduced further when k = n. The derivation of (A.1) uses the estimate

$$||d(u - \tilde{u}_h)||_W \le c(E(du) + \eta[E(d\sigma) + E(p)])$$

from [3, Theorem 2.11] which is unnecessary here since the left side is always zero. Since this is the only part of the derivation that requires the term $\eta E(d\sigma)$, we can drop it, yielding

$$||u - \tilde{u}_h||_{L^2} \le c(E(u) + \eta E(\sigma) + (\delta + \mu)E(d\sigma) + \mu E(P_{\mathfrak{B}}u)).$$
 (A.2)

We now give bounds on each of the terms in (A.2). The coefficients appearing in the abstract estimates can be stated in terms of powers of h in the deRham context. These appear in [3, p. 312] as

$$\eta = O(h), \quad \delta = O(h^{\min(2,r+1)}), \quad \text{and } \mu = O(h^{r+1}).$$

To bound the error terms, Arnold, Falk and Winther define smooth projection operators $\pi_h^k : L^2 \Lambda^k(\Omega) \to \Lambda_h^k$ satisfying optimal convergence rates as stated precisely in [3, Theorem 5.9]. For instance, if Λ_h^k is one of $\mathcal{P}_{r+1}^- \Lambda^k(\mathcal{T}_h)$ or, if $r \ge 1$, $\mathcal{P}_r \Lambda^k(\mathcal{T}_h)$ then

$$\left|\left|w - \pi_h^k w\right|\right|_{L^2\Lambda^k(\Omega)} \le ch^s \left|\left|w\right|\right|_{H^s\Lambda^k(\Omega)}, \quad \text{for } w \in H^s\Lambda^k(\Omega), \quad 0 \le s \le r+1.$$

These types of results bound E(w) in terms of $||w||_{H^s\Lambda^k}$, which is in turn bounded in terms of $||\Delta u||_{H^s\Lambda^k}$ by the regularity hypothesis (6.7). Summarizing these results, we have

$$E(u) \le ch^{s+2} ||\Delta u||_{H^s}$$
$$E(\sigma) \le ch^{s+1} ||\Delta u||_{H^s}$$
$$E(d\sigma) \le ch^s ||\Delta u||_{H^s}$$
$$E(P_{\mathfrak{B}}u) \le ch^{s+2} ||\Delta u||_{H^s}$$

We can now prove (6.11) by collecting results and applying them to (A.2), yielding

$$|\rho(t)||_{L^2} \le c(h^{s+2} + h(h^{s+1}) + (h^{\min(2,r+1)} + h^{r+1})h^s + h^{r+1}h^{s+2}) ||\Delta u(t)||_{H^s}$$

The greatest common factor from the above expression is h^{s+2} hence this is the overall order estimate that can be inferred, as was claimed.

Acknowledgments. MH was supported in part by NSF Awards 0715146, by DOD/DTRA Award HDTRA-09-1-0036, and by NBCR. AG was supported in part by NSF Award 0715146 and by NBCR. YZ was supported in part by NSF DMS 1319110. The authors would like to thank an anonymous referee for pointing out some important subtle properties of the π_h^k operators as well providing various useful suggestions for technical simplifications of the proofs.

REFERENCES

- D. Arnold and H. Chen. Finite element exterior calculus for parabolic problems. arXiv:1209.1142, 2012.
- [2] D. Arnold, R. Falk, and R. Winther. Finite element exterior calculus, homological techniques, and applications. Acta Numerica, pages 1–155, 2006.
- [3] D. Arnold, R. Falk, and R. Winther. Finite element exterior calculus: from Hodge theory to numerical stability. *Bulletin of the American Mathematical Society*, 47(2):281–354, 2010.
- [4] G. Baker. Error estimates for finite element methods for second order hyperbolic equations. SIAM J. Numer. Anal., pages 564–576, 1976.
- [5] R. E. Bank, M. J. Holst, R. Szypowski, and Y. Zhu. Finite element error estimates for critical growth semilinear problems without angle conditions. arXiv:1108.3661, 2011.
- [6] A. Bossavit. Whitney forms: a class of finite elements for three-dimensional computations in electromagnetism. *Proc. of the IEEE*, 135(8):493–500, 1988.
- [7] J. Bramble, A. Schatz, V. Thomée, and L. Wahlbin. Some convergence estimates for semidiscrete Galerkin type approximations for parabolic equations. *SIAM J. Numer. Anal.*, pages 218–241, 1977.
- [8] J. Brüning and M. Lesch. Hilbert complexes. J. Funct. Anal., 108(1):88–132, 1992.
- [9] K. Chrysafinos and L. Hou. Error estimates for semidiscrete finite element approximations of linear and semilinear parabolic equations under minimal regularity assumptions. *SIAM J. Numer. Anal.*, pages 282–306, 2003.
- [10] L. Cowsar, T. Dupont, and M. Wheeler. A priori estimates for mixed finite element methods for the wave equation. *Computer Methods in Applied Mechanics and Engineering*, 82(1-3):205–222, 1990.
- [11] L. Cowsar, T. Dupont, and M. Wheeler. A priori estimates for mixed finite element approximations of second-order hyperbolic equations with absorbing boundary conditions. *SIAM J. Numer. Anal.*, pages 492–504, 1996.
- [12] R. S. Falk and R. Winther. Local bounded cochain projections. *Math. Comp.*, 83(290):2631–2656, 2014.
- [13] R. S. Falk and R. Winther. Double complexes and local cochain projections. Numer. Methods Partial Differential Equations, 31(2):541–551, 2015.

- [14] S. M. Garcia. Improved error estimates for mixed finite-element approximations for nonlinear parabolic equations: the continuous-time case. *Numerical Methods for Partial Differential Equations*, 10(2):129–147, 1994.
- [15] T. Geveci. On the application of mixed finite element methods to the wave equations. *Mathematical Modelling and Numerical Analysis*, 22(2):243–250, 1988.
- [16] Y. Giga. A bound for global solutions of semilinear heat equations. *Communications in Mathematical Physics*, 103(3):415–421, 1986.
- [17] P. W. Gross and P. R. Kotiuga. Electromagnetic theory and computation: a topological approach, volume 48 of Mathematical Sciences Research Institute Publications. Cambridge University Press, Cambridge, 2004.
- [18] D. Henry. Geometric theory of semilinear parabolic equations, volume 61. Springer-Verlag, 1981.
- [19] R. Hiptmair. Finite elements in computational electromagnetism. Acta Numerica, 11:237–339, 2002.
- [20] M. Holst and A. Stern. Geometric variational crimes: Hilbert complexes, finite element exterior calculus, and problems on hypersurfaces. *Found. Comput. Math.*, 12(3):263–293, 2012.
- [21] M. Holst and A. Stern. Semilinear mixed problems on Hilbert complexes and their numerical approximation. *Found. Comput. Math.*, 12(3):363–387, 2012.
- [22] C. Johnson and V. Thomee. Error estimates for some mixed finite element methods for parabolic type problems. *RAIRO Anal. Numér*, 15(1):41–78, 1981.
- [23] A. Lasis and E. Süli. *hp*-version discontinuous Galerkin finite element method for semilinear parabolic problems. *SIAM J. Numer. Anal.*, 45(4):1544–1569, 2007.
- [24] C. Makridakis. On mixed finite element methods for linear elastodynamics. *Numerische Mathematik*, 61(1):235–260, 1992.
- [25] J.-C. Nédélec. Mixed finite elements in \mathbb{R}^3 . Numer. Math., 35(3):315–341, 1980.
- [26] J.-C. Nédélec. A new family of mixed finite elements in \mathbb{R}^3 . Numer. Math., 50(1):57–81, 1986.
- [27] A. Pazy. Semigroups of linear operators and applications to partial differential equations, volume 44 of Applied Mathematical Sciences. Springer, 1983.
- [28] P.-A. Raviart and J. M. Thomas. A mixed finite element method for 2nd order elliptic problems. In Mathematical aspects of finite element methods (Proc. Conf., Consiglio Naz. delle Ricerche (C.N.R.), Rome, 1975), pages 292–315. Lecture Notes in Math., Vol. 606. Springer, Berlin, 1977.
- [29] M. Renardy and R. Rogers. An introduction to partial differential equations, volume 13. Springer Verlag, 2nd edition, 2004.
- [30] B. Ruf and E. Terraneo. The Cauchy problem for a semilinear heat equation with singular initial data. In *Evolution equations, semigroups and functional analysis: in memory of Brunello Terreni*, volume 50. Birkhauser, 2002.
- [31] T. Sun and D. Filippova. Long-time error estimation for semi-linear parabolic equations. *Journal of Computational and Applied Mathematics*, 185(1):1 18, 2006.
- [32] R. Temam. Infinite-dimensional dynamical systems in mechanics and physics, volume 68. Springer verlag, 1997.
- [33] V. Thomée. Galerkin finite element methods for parabolic problems. Springer Verlag, 2006.
- [34] V. Thomée and L. Wahlbin. On Galerkin methods in semilinear parabolic problems. *SIAM J. Numer. Anal.*, pages 378–389, 1975.
- [35] M. Wheeler. A priori L^2 error estimates for Galerkin approximations to parabolic partial differential equations. *SIAM J. Numer. Anal.*, pages 723–759, 1973.
- [36] J. Wrkich and Q. Zhang. Solvability conditions for some semi-linear parabolic equations. *Journal of Differential Equations*, 247(8):2440–2454, 2009.

E-mail address: agillette@math.arizona.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ARIZONA, TUCSON AZ 85721

E-mail address: mholst@math.ucsd.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA SAN DIEGO, LA JOLLA CA 92093

E-mail address: zhuyunr@isu.edu

DEPARTMENT OF MATHEMATICS & STATISTICS, IDAHO STATE UNIVERSITY, POCATELLO ID 83209