

# FINITE ELEMENT EXTERIOR CALCULUS FOR EVOLUTION PROBLEMS

ANDREW GILLETTE AND MICHAEL HOLST

ABSTRACT. Arnold, Falk, and Winther [*Bull. Amer. Math. Soc.* **47** (2010), 281–354] recently showed that mixed variational problems, and their numerical approximation by mixed methods, could be most completely understood using the ideas and tools of *Hilbert complexes*. This led to the development of the Finite Element Exterior Calculus (FEEC) for a large class of linear elliptic problems. More recently, Holst and Stern [arXiv:1005.4455,arXiv:1010.6127] extended the FEEC framework to semi-linear problems, and to problems containing *variational crimes*, allowing for the analysis and numerical approximation of linear and nonlinear geometric elliptic partial differential equations on Riemannian manifolds of arbitrary spatial dimension, generalizing surface finite element approximation theory. In this article, we develop another distinct extension to the FEEC, namely to parabolic and hyperbolic evolution systems, allowing for the treatment of geometric and other evolution problems. Our approach is to combine the recent work on the FEEC for elliptic problems with a classical approach to solving evolution problems via semi-discrete finite element methods, by viewing solutions to the evolution problem as lying in time-parameterized Hilbert spaces (or *Bochner* spaces). Building on classical approaches by Thomée for parabolic problems and Geveci for hyperbolic problems, we establish *a priori* error estimates for Galerkin FEM approximation in the natural Bochner space norms. In particular, we recover the results of Thomée and Geveci for two-dimensional domains and the lowest-order mixed method as a special case, effectively extending their results to arbitrary spatial dimension and to an entire family of mixed methods. We also show how the Holst and Stern framework allows for extensions of these results to certain semi-linear evolution problems.

## CONTENTS

1. Introduction	2
2. Semi-Discrete FEM Error Estimates for Evolution Problems	4
3. The Finite Element Exterior Calculus	6
4. Abstract Evolution Problems and Bochner Spaces	8
5. <i>A Priori</i> Error Estimates for Parabolic Problems	10
6. <i>A Priori</i> Error Estimates for Hyperbolic Problems	15
7. Semi-linear Evolution Problems	19
8. Concluding Remarks	21
9. Acknowledgments	21
Appendix A. Explanation of Semi-Linear Error Estimates	21
References	23

---

*Date:* February 9, 2012.

*Key words and phrases.* FEEC, elliptic equations, evolution equations, nonlinear equations, approximation theory, nonlinear approximation, inf-sup conditions, *a priori* estimates.

MH was supported in part by NSF Awards 0715146, by DOD/DTRA Award HDTRA-09-1-0036, and by NBCR.

AG was supported in part by NSF Award 0715146 and by NBCR.

## 1. INTRODUCTION

More than two decades of research on linear mixed variational problems, and their numerical approximation by mixed methods, recently culminated in the seminal work of Arnold, Falk, and Winther [3]. They showed that these problems could be most completely understood using the ideas and tools of *Hilbert complexes*, leading to the development of the Finite Element Exterior Calculus (FEEC) for elliptic problems. In two related articles [16, 17], Holst and Stern extended the Arnold–Falk–Winther framework to include *variational crimes*, allowing for the analysis and numerical approximation of linear and nonlinear geometric elliptic partial differential equations on Riemannian manifolds of arbitrary spatial dimension, generalizing the existing surface finite element approximation theory in several directions. In the current article, we extend the FEEC in another direction, namely to parabolic and hyperbolic evolution systems. Our approach is to combine the recent work on the FEEC for elliptic problems with a classical approach to solving evolution problems using semi-discrete finite element methods, by viewing solutions to the evolution problem as lying in time-parameterized Hilbert (or *Bochner*) spaces. Building on classical approaches by Thomée for parabolic problems and Geveci for hyperbolic problems, we establish *a priori* error estimates for Galerkin FEM approximation in the natural Bochner space norms. In particular, we recover the results of Thomée and Geveci for two-dimensional domains and the lowest-order mixed method as a special case, effectively extending their results to arbitrary spatial dimension and to an entire family of mixed methods. We also show how the Holst and Stern framework allows for extensions of these results to certain semi-linear evolution problems.

To understand why the finite element exterior calculus (FEEC) has emerged in a natural way to become a major mathematical tool in the development of numerical methods for PDE, consider the vector Laplacean:

$$-\Delta u = -\operatorname{grad} \operatorname{div} u + \operatorname{curl} \operatorname{curl} u,$$

and a natural variational formulation: Find  $u \in H(\operatorname{curl}; \Omega) \cap H_0(\operatorname{div}; \Omega)$  s.t.

$$\int_{\Omega} [(\nabla \cdot u)(\nabla \cdot v) + (\nabla \times u) \cdot (\nabla \times v)] dx = \int_{\Omega} f \cdot v dx, \quad \forall v \in H(\operatorname{curl}; \Omega) \cap H_0(\operatorname{div}; \Omega).$$

A *mixed* formulation is a natural alternative: Find  $(\sigma, u) \in H^1(\Omega) \times H(\operatorname{curl}; \Omega)$  s.t.

$$\int_{\Omega} (\sigma \tau - u \cdot \nabla \tau) dx = 0, \quad \forall \tau \in H^1(\Omega),$$

$$\int_{\Omega} [\nabla \sigma \cdot v + (\nabla \times u) \cdot (\nabla \times v)] dx = \int_{\Omega} f \cdot v dx, \quad \forall v \in H(\operatorname{curl}; \Omega).$$

Using the standard finite element approach based on the first (non-mixed) formulation turns out to not correctly capture either the geometry (in the case of domains with corners) or the topology (in the case of non-simply connected domains); there are several detailed examples in [3] that explore what goes wrong. A standard finite element approach based on the second (mixed) formulation turns out to work extremely well and encounters neither of the difficulties of the first approach. Why is this the case? To answer this question, consider an arbitrary vector field  $f \in L^2(\Omega)$ ; we have the *Helmholtz-Hodge* orthogonal decomposition into curl-free, divergence-free, and harmonic functions:

$$f = \nabla p + \nabla \times q + h,$$

where  $h$  is harmonic (divergence- and curl-free). The mixed formulation is essentially computing this decomposition for  $h = 0$  and finite element methods are somehow exploiting this.

The relationship can be found by examining the *de Rham cohomology* of the domain  $\Omega$ . The space of harmonic  $k$ -forms is isomorphic to the  $k$ th de Rham cohomology group of the domain  $\Omega$ , meaning the dimension of the space of harmonic forms indicates the number of topological features that can create obstacles to well-posed formulations of elliptic problems. A natural question is then: What is an appropriate mathematical framework for understanding this abstractly, that will allow for a methodical construction of “good” finite element methods for these types of problems?

The answer turns out to be *Hilbert Complexes*. Hilbert complexes were originally studied in [8] as a way to generalize certain properties of elliptic complexes, particularly the Hodge decomposition and other aspects of Hodge theory. A *Hilbert complex*  $(W, d)$  consists of a sequence of Hilbert spaces  $W^k$ , along with closed, densely-defined linear maps  $d^k: V^k \subset W^k \rightarrow V^{k+1} \subset W^{k+1}$ , possibly unbounded, such that  $d^k \circ d^{k-1} = 0$  for each  $k$ .

$$\dots \longrightarrow V^{k-1} \xrightarrow{d^{k-1}} V^k \xrightarrow{d^k} V^{k+1} \longrightarrow \dots$$

This Hilbert complex is said to be *bounded* if  $d^k$  is a bounded linear map from  $W^k$  to  $W^{k+1}$  for each  $k$ , i.e.,  $(W, d)$  is a cochain complex in the category of Hilbert spaces. It is said to be *closed* if the image  $d^k V^k$  is closed in  $W^{k+1}$  for each  $k$ . It was shown in [3] that Hilbert complexes are also a convenient abstract setting for mixed variational problems and their numerical approximation by mixed finite element methods, providing the foundation of a framework called *finite element exterior calculus* (see also [2]). This line of research is the culmination of several decades of work on mixed finite element methods and computational electromagnetics [6, 20, 21, 14]. The most important example of a Hilbert complex for our purposes of the FEEC arises from the de Rham complex of smooth differential forms on a domain or manifold.

The main developments in FEEC to date have been for linear (and now semi-linear) elliptic problems such as Poisson’s equation

$$-\Delta u = f.$$

Our goal here is expand the scope of this analysis to include parabolic linear (and semi-linear) equations such as the heat equation,

$$(\partial_t - \Delta)u = f,$$

and hyperbolic equations such as the wave equation,

$$(\partial_{tt} - \Delta)u = f.$$

The exterior calculus framework treats  $\Delta$  as  $(d + \delta)^2$ , where  $d$  is the exterior derivative operator and  $\delta$  its adjoint. The incorporation of the time derivative operation  $\partial_t$  into this framework, however, has not been previously considered. To remedy this, we develop the most natural extension of FEEC theory to evolution problems: a generalization of the semi-discrete method often called the ‘method of lines.’ This approach involves the discretization of the spatial part of the differential operator, leaving the time variable continuous. It can be viewed as introducing a time parameter into the discrete (Hilbert) spaces that have been developed for elliptic problems. These parametrized Hilbert spaces are naturally treated as *Bochner spaces* and we will show how Bochner space theory makes obvious the well-posedness of the problems we consider. Moreover, the accompanying Bochner space norms, when coupled with FEEC notation for Hilbert complexes, provide

a clear and consistent notation for bounding errors in mixed methods accumulated over a finite time interval.

We note that there is another approach to solving evolution problems with finite elements, namely using a complete discretization of space-time. This tactic allows for the dynamical change of the underlying discrete approximation spaces in both space and time. Such an approach gives rise to space-time adaptivity, and is potentially the most flexible and powerful approach to the numerical treatment of parabolic and hyperbolic evolution problems. This approach, which we will consider in a second article, is most naturally formulated using geometric calculus, a well-studied mathematical structure for time-dependent problems. In the current article, we focus on extending FEED to semi-discrete methods using Bochner norm estimates for the method-of-lines approach.

Finally, we note that the work presented here was developed simultaneously and independently from a related project by Arnold and Chen [1]. While our focus is extending the *scalar* Hodge-Laplacian to both linear and semi-linear evolution problems, with or without geometric variational crimes, their focus is on extending FEED theory to generalized Hodge-Laplacian parabolic problems. The pairing of these two results will lead to further insight on the subject in a variety of directions.

*Outline of the paper.* The remainder of the paper is structured as follows. In Section 2, we review the classical semi-discrete mixed finite element method error estimates for parabolic problems due to Thomée [27] and others for hyperbolic problems due to Geveci [12]. In Section 3, we give a very brief overview the Finite Element Exterior Calculus and recall some relevant results. In Section 4, we formulate abstract parabolic and hyperbolic problems in Bochner spaces and state some standard results on the existence and uniqueness of strong and weak solutions. In Section 5, we combine the classical approach to semi-discrete methods with modern FEED theory to establish some basic *a priori* error estimates for Galerkin mixed finite element methods for parabolic problems. The main result is Theorem 5.2, which exploits the FEED framework to obtain a classification of spatial finite element spaces that give optimal order convergence rates in Bochner norms. In Section 6, we carry out a similar analysis for hyperbolic problems, resulting in the error estimate given in Theorem 6.2. Our results recover the estimates of Thomée and Geveci for two-dimensional domains and the lowest-order mixed method as a special case, effectively extending their results to arbitrary spatial dimension and to an entire family of mixed methods. In Section 7, we employ the results of Holst and Stern to extend our parabolic estimates to a class of semi-linear evolution PDE. Finally, in Section 8, we draw some conclusions and make some remarks on future directions.

## 2. SEMI-DISCRETE FEM ERROR ESTIMATES FOR EVOLUTION PROBLEMS

We begin by reviewing semi-discrete finite element methods and their *a priori* error estimates for parabolic and hyperbolic PDE systems. We focus in each case on a relatively simple, well-studied system of interest to modeling communities, namely, the heat equation (parabolic) and the wave equation (hyperbolic). The **heat equation** is: find  $u(x, t)$  such that

$$\begin{aligned} u_t - \Delta u &= f && \text{in } \Omega, && \text{for } t > 0 \\ u &= 0 && \text{on } \partial\Omega, && \text{for } t > 0 \quad \text{with } u(\cdot, 0) = g \text{ in } \Omega. \end{aligned} \tag{2.1}$$

We review the approach to Galerkin methods for this problem as presented in Thomée [27] for domains  $\Omega \subset \mathbb{R}^2$ . His approach is based on an earlier work with Johnson [18] and builds upon prior analysis of elliptic projection [7]. Let  $\sigma = \nabla u$  and define the mixed,

weak form problem: Find  $(u, \sigma) \in L^2 \times H(\text{div})$  such that

$$\begin{aligned} (u_t, \phi) - (\text{div} \sigma, \phi) &= (f, \phi), \quad \forall \phi \in L^2, \quad t > 0 \\ (\sigma, \omega) + (u, \text{div} \omega) &= 0, \quad \forall \omega \in H(\text{div}), \quad t > 0, \quad u(0) = g. \end{aligned} \quad (2.2)$$

The semi-discrete problem is then to find  $(u_h, \sigma_h) \in S_h \times H_h \subset L^2 \times H(\text{div})$  such that

$$\begin{aligned} (u_{h,t}, \phi_h) - (\text{div} \sigma_h, \phi_h) &= (f, \phi_h), \quad \forall \phi_h \in S_h, \quad t > 0 \\ (\sigma_h, \omega_h) + (u_h, \text{div} \omega_h) &= 0, \quad \forall \omega_h \in H_h, \quad t > 0, \quad u_h(0) = g_h. \end{aligned} \quad (2.3)$$

where  $g_h$  is an approximation of  $g$  in  $S_h$ . With bases for  $S_h$  and  $H_h$ , the matrix form of the discrete problem is

$$\begin{aligned} AU_t - B\Sigma &= F, \\ B^T U + D\Sigma &= 0, \quad \text{for } t > 0, \quad U(0) \text{ given,} \end{aligned}$$

where  $U$  and  $\Sigma$  are vectors corresponding to  $u_h$  and  $\sigma_h$ . It is easily seen that the matrices  $A$  and  $D$  are positive definite. Eliminating  $\Sigma$ , we have the system of ODEs

$$AU_t + BD^{-1}B^T U = F, \quad \text{for } t > 0, \quad U(0) \text{ given,}$$

which by standard results in ODE theory has a unique solution.

Thomée uses discontinuous linear elements for  $S_h$  and piecewise quadratic elements for  $H_h$ . He defines the solution operator  $T_h : L^2 \rightarrow S_h$  given by  $T_h f = u_h$  for the corresponding elliptic problem. Then  $g_h$  can be defined by

$$g_h := R_h g := -T_h \Delta g.$$

For  $g_h = R_h g$  and  $t \geq 0$ , Thomée derives the estimates

$$\|u_h(t) - u(t)\|_{L^2} \leq ch^2 \left( \|u(t)\|_{H^2} + \int_0^t \|u_t\|_{H^2} ds \right), \quad (2.4)$$

$$\|\sigma_h(t) - \sigma(t)\|_{L^2} \leq ch^2 \left( \|u(t)\|_{H^3} + \left( \int_0^t \|u_t\|_{H^2}^2 ds \right)^{1/2} \right). \quad (2.5)$$

Note that these estimates are for a fixed time value  $t$  and restricted to a particular choice of finite elements in 2D.

We now turn to the **wave equation**: find  $u(x, t)$  such that

$$\begin{aligned} u_{tt} - \Delta u &= f \quad \text{in } \Omega, \quad \text{for } t > 0, \\ u &= 0 \quad \text{on } \partial\Omega, \quad \text{for } t > 0 \quad \text{with } u(\cdot, 0) = u_0 \text{ in } \Omega, \\ & \quad \text{and } u_t(\cdot, 0) = u_1 \text{ in } \Omega \end{aligned} \quad (2.6)$$

There are two approaches to defining a mixed weak form of this problem. The first is very similar to the parabolic case: given  $f$ ,  $u_0$ , and  $u_1$ , find  $(u, \sigma)$  such that

$$\begin{aligned} (u_{tt}, \phi) - (\text{div} \sigma, \phi) &= (f, \phi), \quad \forall \phi \in L^2, \quad t > 0, \\ (\sigma, \omega) + (u, \text{div} \omega) &= 0, \quad \forall \omega \in H(\text{div}), \quad t > 0, \\ u(0) &= u_0, \\ u_t(0) &= u_1. \end{aligned} \quad (2.7)$$

It is difficult to derive estimates for the numerical approximation of (2.7) akin to those found in the parabolic case due to the second derivatives appearing in the formulation. Some estimates attempts along these lines for  $\Omega \subset \mathbb{R}^2$  have been given by Baker [4] and Cowsar, Dupont and Wheeler [10, 11].

For the purpose of extending the FEEC framework, we find the **‘velocity-stress’ formulation** of the problem and the results of Geveci [12] to be more useful. This formulation solves for  $\mu := u_t$  instead of  $u$ : given  $f$ ,  $u_0$ , and  $u_1$ , find  $(\mu, \sigma) \in L^2 \times H(\text{div})$  such that

$$\begin{aligned} (\mu_t, \phi) - (\text{div } \sigma, \phi) &= (f, \phi), \quad \forall \phi \in L^2, \quad t > 0, \\ (\sigma_t, \omega) + (\mu, \text{div } \omega) &= 0, \quad \forall \omega \in H(\text{div}), \quad t > 0, \\ \mu(0) &= u_1, \\ \sigma(0) &= \nabla u_0. \end{aligned} \tag{2.8}$$

The semi-discrete problem is then to find  $(\mu_h, \sigma_h) \in S_h \times H_h \subset L^2 \times H(\text{div})$  such that

$$\begin{aligned} (\mu_{h,t}, \phi_h) - (\text{div } \sigma_h, \phi_h) &= (f, \phi_h), \quad \forall \phi_h \in S_h, \quad t > 0, \\ (\sigma_{h,t}, \omega_h) + (\mu_h, \text{div } \omega_h) &= 0, \quad \forall \omega_h \in H_h, \quad t > 0, \\ \mu_h(0) &= u_{1,h}, \\ \sigma_h(0) &= (\nabla u_0)_h, \end{aligned} \tag{2.9}$$

where  $u_{1,h}$  is an approximation of  $u_1$  in  $S_h$  and  $(\nabla u_0)_h$  is an approximation of  $\nabla u_0$ . Again, bases for  $S_h$  and  $H_h$  reduce the discrete problem to a matrix formulation:

$$\begin{aligned} AW_t - B\Sigma &= F, \\ B^T W + D\Sigma_t &= 0, \quad \text{for } t > 0, \quad W(0), \Sigma(0) \text{ given,} \end{aligned}$$

where  $W$  and  $\Sigma$  are vectors corresponding to  $\mu_h$  and  $\sigma_h$  and  $A, D$  are symmetric, positive definite matrices. As Geveci [12, p. 248] explains, this can be reduced to a single iterative system of the form

$$(D + k^2 B^T A^{-1} B) \Sigma^{n+1} = G,$$

where  $k$  denotes the time step in an implicit Euler time-differencing scheme.

To derive an error estimate for the velocity-stress discretization, Geveci states the need for projection operators from  $H(\text{div})$  to  $H_h$  and from  $L^2$  to  $S_h$  satisfying certain approximation properties. He explains that such operators exist for a variety of finite element spaces in  $\mathbb{R}^2$ , e.g. the Raviart-Thomas spaces [22], allowing the following result. For  $1 \leq s \leq r$  with  $r \geq 2$ ,

$$\begin{aligned} \|\mu_h(t) - \mu(t)\|_{L^2} + \|\sigma_h(t) - \sigma(t)\|_{L^2} &\leq c(\|u_1 - u_{1,h}\| + \|\nabla u_0 - (\nabla u_0)_h\|) + \\ &+ ch^s \left( \|u_1\|_s + \|\nabla u_0\|_s + \int_0^t (\|\mu_t(\tau)\|_s + \|\sigma_t(\tau)\|_s) d\tau \right). \end{aligned} \tag{2.10}$$

Like estimates (2.4) and (2.5) for the parabolic problem, (2.10) says that the approximation error can be controlled in  $L^2$  norm at any time  $t$  by the  $H^s$  norm of the initial conditions plus the accumulated norm of the variables up to time  $t$ . It is these types of estimates that the FEEC framework can refine, simplify, and generalize to arbitrary spatial dimension  $n$ .

### 3. THE FINITE ELEMENT EXTERIOR CALCULUS

The finite element exterior calculus (FEEC) provides an elegant mathematical framework for deriving error estimates for a large class of elliptic PDE. We now give a brief overview of the notation and certain main results from FEEC which are relevant to this paper; we refer the reader to the seminal papers of Arnold, Falk, and Winther [2, 3] for additional explanation.

Let  $\Omega$  be a bounded  $n$ -manifold embedded in  $\mathbb{R}^n$  and assume  $\Omega$  has a piecewise smooth, Lipschitz boundary. The space of  $L^2$ -bounded continuous differential  $k$ -forms on  $\Omega$  is given by

$$L^2\Lambda^k(\Omega) := \left\{ \sum_I a_I dx_I \in \Lambda^k(\Omega) : a_I \in L^2(\Omega) \quad \forall I \right\},$$

where  $I$  ranges over all strictly increasing sequences of  $k$  indices chosen from  $\{1, \dots, n\}$ . The exterior derivative operator  $d_k : \Lambda^k(\Omega) \rightarrow \Lambda^{k+1}(\Omega)$  acts on these spaces to form a Hilbert complex  $(L^2\Lambda, d)$ . The associated domain complex is the sequence of spaces  $H\Lambda^k := \text{domain}(d_k) \subset L^2\Lambda^k(\Omega)$ , commonly called the  $L^2$  deRham complex:

$$0 \longrightarrow H\Lambda^0 \xrightarrow{d_0} H\Lambda^1 \xrightarrow{d_1} \dots \xrightarrow{d_{n-1}} H\Lambda^n \longrightarrow 0.$$

The norm on each space is the graph norm associated to  $d$ , i.e.

$$(u, v)_{H^k(\Omega)} := (u, v)_{\Lambda^k(\Omega)} + (d_k u, d_k v)_{\Lambda^{k+1}(\Omega)}.$$

We note that in any dimension  $n$ , the beginning and end of the  $L^2$  deRham complex can be understood in terms of traditional Sobolev spaces and differential operators:

$$0 \longrightarrow H^1(\Omega) \xrightarrow{\text{grad}} \dots \xrightarrow{\text{div}} L^2(\Omega) \longrightarrow 0$$

A major conclusion of FEEC is that stable finite element methods for elliptic PDE must seek solutions in finite dimensional subspaces  $\Lambda_h^k \subset H\Lambda^k$  that satisfy certain key approximation properties. First, the subspaces should form a subcomplex of the  $L^2$  deRham complex, meaning  $d\Lambda_h^k \subset \Lambda_h^{k+1}$ . Second,  $\Lambda_h^k$  should have sufficient approximation that upper bounds on  $\inf_{v \in \Lambda_h^k} \|u - v\|_{H\Lambda^k}$  can be ensured for some or all  $u \in H\Lambda^k$ . Third, there must exist bounded cochain projections  $\pi_h^k : H\Lambda^k \rightarrow \Lambda_h^k$  which are invariant on  $\Lambda_h^k$ , commute with the exterior derivative operators, and provide a bound  $\|\pi_h^k v\|_{H\Lambda^k} \leq c \|v\|_{H\Lambda^k}$  for all  $v \in H\Lambda^k$ .

In the context of the deRham complex, all these properties are shown to be provided for by two canonical classes of piecewise degree  $r$  polynomials associated to a simplicial mesh  $\mathcal{T}$  of  $\Omega$ . Let  $\mathcal{P}_r$  denote polynomials in  $n$  variables of degree at most  $r$  and  $\mathcal{H}_r \subset \mathcal{P}_r$  the subspace of homogeneous polynomials. The first class, denoted  $\mathcal{P}_r\Lambda^k(\mathcal{T})$ , consists of all  $k$ -forms with coefficients belonging to  $\mathcal{P}_r$  on each  $n$ -simplex of  $\mathcal{T}$ . The second class, denoted  $\mathcal{P}_r^-\Lambda^k(\mathcal{T})$ , interleaves with the first class, i.e.

$$\mathcal{P}_{r-1}\Lambda^k(\mathcal{T}) \subsetneq \mathcal{P}_r^-\Lambda^k(\mathcal{T}) \subsetneq \mathcal{P}_r\Lambda^k(\mathcal{T}).$$

To define  $\mathcal{P}_r^-\Lambda^k(\mathcal{T})$ , first define  $X$  be the vector field on  $\mathbb{R}^n$  such that  $X(x)$  is the vector based at  $x \in \mathbb{R}^n$  that points opposite to the origin with length  $|x|$ . Define  $\mathcal{P}_r^-\Lambda^k(\mathcal{T}) := \mathcal{P}_r\Lambda^k \oplus \kappa\mathcal{H}_{r-1}\Lambda^{k+1}$ , a direct sum, where  $\kappa$  is defined by contraction with  $X$ . The map  $\kappa$  is called the *Koszul differential* and gives rise to the *Koszul complex*. This is elaborated upon in detail in the work of Arnold, Falk and Winther e.g. [3, p. 328].

For  $n = 3$ , we have the following correspondences between the FEEC notation of finite element spaces and traditional element spaces.

$$\begin{aligned} \mathcal{P}_{r+1}\Lambda^2(\mathcal{T}) &= \text{Nédélec 2nd-kind } H(\text{div}) \text{ elements of degree } \leq r + 1 \text{ (see [21])} \\ \mathcal{P}_{r+1}^-\Lambda^2(\mathcal{T}) &= \text{Nédélec 1st-kind } H(\text{div}) \text{ elements of order } r \text{ (see [20])} \\ \mathcal{P}_{r+1}^-\Lambda^3(\mathcal{T}) &= \text{discontinuous elements of degree } \leq r \\ \mathcal{P}_r\Lambda^3(\mathcal{T}) &= \text{discontinuous elements of degree } \leq r \end{aligned}$$

Hence, in the case of the deRham complex, FEEC recovers the well-known finite element spaces while at the same time describing their generalization to arbitrary spatial dimensions.

The last piece of FEEC used in this work is the existence of smoothed projection operators

$$\pi_h^k : L^2 \Lambda^k \rightarrow \Lambda_h^k \quad \text{where } \Lambda_h^k \in \{\mathcal{P}_r \Lambda^k(\mathcal{T}), \mathcal{P}_r^- \Lambda^k(\mathcal{T})\}. \quad (3.1)$$

The following theorem asserts the key properties of these operators.

**Theorem 3.1** ([3] Theorem 5.9).

(1) Let  $\Lambda_h^k$  be one of the spaces  $\mathcal{P}_{r+1}^- \Lambda^k(\mathcal{T})$  or, if  $r \geq 1$ ,  $\mathcal{P}_r \Lambda^k(\mathcal{T})$ . Then  $\pi_h^k$  is a projection onto  $\Lambda_h^k$  and satisfies

$$\|\omega - \pi_h^k \omega\|_{L^2 \Lambda^k(\Omega)} \leq ch^s \|\omega\|_{H^s \Lambda^k(\Omega)}, \quad \omega \in H^s \Lambda^k(\Omega),$$

for  $0 \leq s \leq r + 1$ . Moreover, for all  $\omega \in L^2 \Lambda^k(\Omega)$ ,  $\pi_h^k \omega \rightarrow \omega$  in  $L^2$  as  $h \rightarrow 0$ .

(2) Let  $\Lambda_h^k$  be one of the spaces  $\mathcal{P}_r \Lambda^k(\mathcal{T})$  or  $\mathcal{P}_r^- \Lambda^k(\mathcal{T})$  with  $r \geq 1$ . Then

$$\|d(\omega - \pi_h^k \omega)\|_{L^2 \Lambda^k(\Omega)} \leq ch^s \|d\omega\|_{H^s \Lambda^k(\Omega)}, \quad \omega \in H^s \Lambda^k(\Omega),$$

for  $0 \leq s \leq r$ .

(3) Let  $\Lambda_h^{k-1} \in \{\mathcal{P}_{r+1} \Lambda^{k-1}(\mathcal{T}), \mathcal{P}_{r+1}^- \Lambda^{k-1}(\mathcal{T})\}$  and  $\Lambda_h^k = \mathcal{P}_{r+1}^- \Lambda^k(\mathcal{T})$  or, if  $r > 0$ ,  $\mathcal{P}_r \Lambda^k(\mathcal{T})$ . Then  $d\pi_h^{k-1} = \pi_h^k d$ .

An explicit construction of these operators can be found in the papers of Arnold, Falk, and Winther [2, 3].

#### 4. ABSTRACT EVOLUTION PROBLEMS AND BOCHNER SPACES

We now cast parabolic and hyperbolic problems into the abstract framework of Bochner spaces, i.e. spaces of functions whose values lie in a Banach space. This approach to analyzing parabolic and hyperbolic evolution PDE can be found for example in [26, page 66] and [23].

Let  $X$  be a Banach space and  $\mathring{I} := (0, T)$  an interval with closure  $\bar{I} := [0, T]$ . Define

$$C(\mathring{I}, X) := \{u : \mathring{I} \rightarrow X \mid u \text{ bounded and continuous}\}.$$

Equip this space with the norm

$$\|u\|_{C(\mathring{I}, X)} := \sup_{t \in \mathring{I}} \|u(t)\|_X.$$

The Bochner space  $L^p(\mathring{I}, X)$  is then defined to be the completion of  $C(\mathring{I}, X)$  with respect to the norm

$$\|u\|_{L^p(\mathring{I}, X)} := \left( \int_{\mathring{I}} \|u(t)\|_X^p dt \right)^{1/p}.$$

The space  $H^1(\mathring{I}, X)$  has an analogous norm

$$\|u\|_{H^1(\mathring{I}, X)} := \left( \int_{\mathring{I}} \|u(t)\|_X^2 + \left\| \frac{d}{dt} u(t) \right\|_X^2 dt \right)^{1/2}.$$

We now consider abstractions of the two main types of evolution PDE.

Let  $H$  and  $V$  be real, separable Banach spaces such that  $V$  is continuously and densely embedded in  $H$ . This provides a Gelfand triple, also called a ‘rigged Hilbert space’:

$$V \subset H \subset V^*,$$

where  $H$  is also continuously and densely embedded in  $V^*$ . It should be noted that the isomorphism between  $V$  and  $V^*$  is in general *not* the same as the composition of the inclusion mappings.

Let  $(\cdot, \cdot)$  denote the inner product on  $H$  as well as the natural pairing between  $V^*$  and  $V$ . Given  $A(t) \in \mathcal{L}(V, V^*)$  depending continuously on  $t \in \bar{I}$ , define a quadratic form

$$a(t, u, v) := -(A(t)u, v), \quad (4.1)$$

for  $(t, u, v) \in \mathbb{R} \times V \times V$ . Assume that  $a$  satisfies the coercivity condition

$$a(t, u, u) \geq c_1 \|u\|_V^2 - c_2 \|u\|_H^2, \quad (4.2)$$

with  $c_1, c_2$  constants independent of  $t \in \bar{I}$ . Consider the abstract parabolic problem

$$u_t = A(t)u + f(t), \quad (4.3)$$

$$u(0) = u_0, \quad (4.4)$$

and the abstract hyperbolic problem

$$\ddot{u} = A(t)u + f(t), \quad (4.5)$$

$$u(0) = u_0, \quad (4.6)$$

$$\dot{u}(0) = u_1. \quad (4.7)$$

These abstract formulations are well-posed in the following sense.

**Theorem 4.1** (Existence of Unique Solution to the Abstract Parabolic Problem). *Let  $f \in L^2(I, V^*)$  and  $u_0 \in H$ , let  $a(\cdot, \cdot, \cdot)$  be as in (4.1), and let (4.2) hold. Then the abstract parabolic problem (4.3) has a unique solution*

$$u \in L^2(\overset{\circ}{I}, V) \cap H^1(\overset{\circ}{I}, V^*).$$

*Moreover, the Sobolev embedding theorem implies  $u \in C(\bar{I}, V^*)$ , allowing an interpretation of the initial condition  $u(0) = u_0$ .*

*Proof.* See [23], page 382. □

The analogous result for the abstract hyperbolic case will require two additional conditions:

$$a(t, u, v) = a(t, v, u), \quad \forall u, v \in V, \quad (4.8)$$

$$A \in C^1(\bar{I}, \mathcal{L}(V, V^*)). \quad (4.9)$$

We now have

**Theorem 4.2** (Existence of Unique Weak Solution to the Abstract Hyperbolic Problem). *Given  $f \in L^1(I, H)$ ,  $u_0 \in V$ , and  $u_1 \in H$ . Let  $a(\cdot, \cdot, \cdot)$  be as in (4.1), and let (4.2), (4.8), and (4.9) hold. Then the abstract hyperbolic problem (4.5) has a unique weak solution*

$$u \in C(\bar{I}, V) \cap C^1(\bar{I}, H).$$

*Proof.* See [23], page 389. □

These results give a short and elegant way to prove that a wide class of PDE problems amenable to finite element methods are well-posed.

### 5. *A Priori* ERROR ESTIMATES FOR PARABOLIC PROBLEMS

We extend Thomée's error estimates from Section 2 to the broader class of elements and arbitrary spatial dimension allowed by FEEC using the abstract framework established in Section 4. Fix an interval  $I = [0, T]$  and define the  $L^2$  Bochner differential  $k$ -form space by parametrizing the  $L^2$  deRham complex and its domain complex. For these spaces, we use the notation

$$L^2 \mathfrak{X}^k := L^2(I, L^2 \Lambda^k(\Omega)) \quad \text{and} \quad H\mathfrak{X}^k := L^2(I, H\Lambda^k(\Omega)). \quad (5.1)$$

We first observe that the parabolic problem (2.1) is a special case of the abstract parabolic problem (4.3)-(4.4). Thus by Theorem 4.1, it has a unique solution on  $(0, T)$  with a  $C^0$ -continuous extension to  $[0, T]$ . We may therefore denote the interval of solution to be  $I := [0, T]$  without confusion. For  $\Omega \subset \mathbb{R}^n$ , define the **Bochner mixed weak parabolic problem**: Given  $f$  and  $g$ , find  $(u, \sigma) \in H\mathfrak{X}^n \times H\mathfrak{X}^{n-1}$  such that

$$\begin{aligned} (u_t, \phi) - (\operatorname{div} \sigma, \phi) &= (f, \phi), \quad \forall \phi \in H\Lambda^n, \quad t \in I, \\ (\sigma, \omega) + (u, \operatorname{div} \omega) &= 0, \quad \forall \omega \in H\Lambda^{n-1}, \quad t \in I, \\ u(0) &= g. \end{aligned} \quad (5.2)$$

The **semi-discrete Bochner parabolic problem** is: Find  $(u_h, \sigma_h) \in \Lambda_h^n \times \Lambda_h^{n-1}$  such that

$$\begin{aligned} (u_{h,t}, \phi_h) - (\operatorname{div} \sigma_h, \phi_h) &= (f, \phi_h), \quad \forall \phi_h \in \Lambda_h^n, \quad t \in I, \\ (\sigma_h, \omega_h) + (u_h, \operatorname{div} \omega_h) &= 0, \quad \forall \omega_h \in \Lambda_h^{n-1}, \quad t \in I, \\ u_h(0) &= g_h. \end{aligned} \quad (5.3)$$

where  $g_h \in \Lambda_h^n$  is an approximation of  $g$ . Define  $g_h$  to be the solution to the elliptic problem with load data  $-\Delta g$ , i.e.

$$\begin{aligned} (\operatorname{div} \hat{\sigma}_h, \phi_h) + (\Delta g, \phi_h) &= 0 \quad \forall \phi_h \in \Lambda_h^n, \\ (\hat{\sigma}_h, \omega_h) + (g_h, \operatorname{div} \omega_h) &= 0, \quad \forall \omega_h \in \Lambda_h^{n-1}. \end{aligned} \quad (5.4)$$

Viewed another way,  $g_h$  is the image of the solution operator  $T_h : L^2 \rightarrow \Lambda_h^n$  for the elliptic problem, i.e.  $g_h = -T_h \Delta g$ . It can be shown, in the same manner presented in [27], that  $T_h$  is positive definite on  $\Lambda_h^n$  implying that the semi-discrete Bochner problem has a unique solution.

Elliptic projection, an idea dating back to Wheeler [29], can be carried out for any fixed time value as we now discuss. For any  $t_0 \in I$ , define the **time-ignorant discrete elliptic problem**: find  $(\tilde{u}_h, \tilde{\sigma}_h) \in \Lambda_h^n \times \Lambda_h^{n-1}$  such that

$$\begin{aligned} (\operatorname{div} \tilde{\sigma}_h, \phi_h) + (-\Delta u(t_0), \phi_h) &= 0, \quad \forall \phi_h \in \Lambda_h^n, \\ (\tilde{\sigma}_h, \omega_h) + (\tilde{u}_h, \operatorname{div} \omega_h) &= 0, \quad \forall \omega_h \in \Lambda_h^{n-1}, \\ \tilde{u}_h(0) &= g_h. \end{aligned} \quad (5.5)$$

Note that the  $u$  appearing in the first equation of (5.5) is the solution to the continuous problem (5.2). Thus, we can view  $\tilde{\sigma}_h$  and  $\tilde{u}_h$  as functions of  $t$  with the understanding that they are defined for each  $t$  value by (5.5) alone; no continuity with respect to  $t$  is required, hence the moniker 'time-ignorant.'

For ease of notation, and in keeping with Thomée, define the error functions

$$\begin{aligned} \rho(t) &:= \tilde{u}_h(t) - u(t), \\ \theta(t) &:= u_h(t) - \tilde{u}_h(t), \\ \varepsilon(t) &:= \sigma_h(t) - \tilde{\sigma}_h(t). \end{aligned}$$

We now prove a lemma which will aid in our subsequent analysis; the result for the linear parabolic problem appears as part of the proof of Thomée [27, Theorem 17.2].

**Lemma 5.1** (Thomé [27]). *The error functions satisfy the semi-discrete formulation:*

$$\begin{aligned} (\theta_t, \phi_h) - (\operatorname{div} \varepsilon, \phi_h) &= -(\rho_t, \phi_h), \quad \forall \phi_h \in \Lambda_h^n, \quad t \in I, \\ (\varepsilon, \omega_h) + (\theta, \operatorname{div} \omega_h) &= 0, \quad \forall \omega_h \in \Lambda_h^{n-1}, \quad t \in I. \end{aligned} \quad (5.6)$$

*Proof.* The second equation is immediate from the second equations in (5.3) and (5.5). The first equation can be written out as

$$(u_{h,t}, \phi_h) - (\operatorname{div} \sigma_h, \phi_h) + (\operatorname{div} \tilde{\sigma}_h, \phi_h) - (\tilde{u}_{h,t}, \phi_h) = (u_t, \phi_h) - (\tilde{u}_{h,t}, \phi_h)$$

which is reduced as follows:

$$\begin{aligned} (u_{h,t}, \phi_h) - (\operatorname{div} \sigma_h, \phi_h) + (\operatorname{div} \tilde{\sigma}_h, \phi_h) &= (u_t, \phi_h) && \text{cancel like terms} \\ (u_{h,t}, \phi_h) - (\operatorname{div} \sigma_h, \phi_h) &= -(\Delta u, \phi_h) + (u_t, \phi_h) && \text{by (5.5)} \\ (f, \phi_h) &= -(\Delta u, \phi_h) + (u_t, \phi_h) && \text{by (5.3)} \end{aligned}$$

This says that the continuous problem  $u_t - \Delta u = f$  should hold in a weak sense when tested against any of the functions in  $\Lambda_h^n$ . This is guaranteed to be true since we chose  $\Lambda_h^n \subset \Lambda^n = L^2$ . Thus, the error equations hold as stated.  $\square$

The following theorem says that if  $\Lambda_h^n$  and  $\Lambda_h^{n-1}$  are chosen according to the FEEC framework, then error estimates akin to Thomée's can be obtained.

**Theorem 5.2.** *Fix  $I := [0, T]$ . Suppose  $(u, \sigma)$  is the solution to (5.2) such that the regularity estimate <sup>1</sup>*

$$\|u(t)\|_{H^{s+2}} + \|du(t)\|_{H^{s+1}} + \|\sigma(t)\|_{H^{s+1}} + \|d\sigma(t)\|_{H^s} \leq c \|f(t)\|_{H^s} \quad (5.7)$$

*holds for  $0 \leq s \leq s_{\max}$  and  $t \in I$ . Choose finite element spaces*

$$\Lambda_h^{n-1} = \left\{ \begin{array}{c} \mathcal{P}_{r+1} \Lambda^{n-1}(\mathcal{T}) \\ \text{or} \\ \mathcal{P}_{r+1}^- \Lambda^{n-1}(\mathcal{T}) \end{array} \right\}, \quad \Lambda_h^n = \mathcal{P}_{r+1}^- \Lambda^n(\mathcal{T}) \quad (= \mathcal{P}_r \Lambda^n(\mathcal{T}))$$

*Then for  $0 \leq s \leq s_{\max}$ ,  $g_h$  defined by (5.4), and  $(u_h, \sigma_h)$  the solution to (5.3), the following error estimates hold:*

$$\|u_h - u\|_{L^2 \mathfrak{X}^n} \leq \begin{cases} ch \left( \|f\|_{L^2(I, L^2)} + \sqrt{T} \|f_t\|_{L^1(I, L^2)} \right) & \text{if } r = 0 \\ ch^{2+s} \left( \|f\|_{L^2(I, H^s)} + \sqrt{T} \|f_t\|_{L^1(I, H^s)} \right) & \text{for } r > 0, \text{ if } s \leq r - 1 \end{cases} \quad (5.8)$$

<sup>1</sup>The regularity assumption (5.7) comes from Arnold, Falk, and Winther [3, p. 342]. Note that  $du = 0$  since  $u \in \Lambda^n$ ; we leave the associated term only to show consistency with [3].

$$\|\sigma_h - \sigma\|_{L^2 \mathfrak{X}^{n-1}} \leq \begin{cases} ch \left( \|f\|_{L^2(I, H^s)} + \sqrt{T} \|f_t\|_{L^2(I, L^2)} \right) \\ \quad \text{if } r = 0, s = 0, \Lambda_h^{n-1} = \mathcal{P}_1^- \Lambda^{n-1}(\mathcal{T}) \\ c \left( h^{1+s} \|f\|_{L^2(I, H^s)} + h\sqrt{T} \|f_t\|_{L^2(I, L^2)} \right) \\ \quad \text{if } r = 0, s \leq 1, \Lambda_h^{n-1} = \mathcal{P}_1 \Lambda^{n-1}(\mathcal{T}) \\ c \left( h^{1+s} \|f\|_{L^2(I, H^s)} + h^{(3/2)+s} \sqrt{T} \|f_t\|_{L^2(I, H^s)} \right) \\ \quad \text{for } r > 0, \text{ if } s \leq r - 1 \end{cases} \quad (5.9)$$

$$\|\operatorname{div}(\sigma_h - \sigma)\|_{L^2 \mathfrak{X}^n} \leq \begin{cases} c \left( h^s \|f\|_{L^2(I, H^s)} + h \|f_t\|_{L^2(I, L^2)} \right) \\ \quad \text{if } r = 0, s \leq 1 \\ c \left( h^s \|f\|_{L^2(I, H^s)} + h^{2+s} \|f_t\|_{L^2(I, H^s)} \right) \\ \quad \text{for } r > 0, \text{ if } s \leq r - 1 \end{cases} \quad (5.10)$$

*Proof.* To simplify notation, we will often use  $\|\cdot\|$  to mean  $\|\cdot\|_{L^2(\Omega)}$ . We adapt the proof technique of the corresponding theorem from Thomée [27, Theorem 17.2] to our setting.

Observe that (5.5) is exactly the  $k = n$  case of the Hodge-Laplacian problem analyzed by Arnold, Falk and Winther [3] and the hypotheses here match their hypotheses. We can thus use a triangle inequality argument for each estimate, e.g.

$$\|u(t) - u_h(t)\|_{L^2} \leq \|u(t) - \tilde{u}_h(t)\|_{L^2} + \|\tilde{u}_h(t) - u_h(t)\|_{L^2} = \|\rho(t)\|_{L^2} + \|\theta(t)\|_{L^2} \quad (5.11)$$

The first term will be bounded using the estimates from [3] and the second by the techniques from Thomée [27]. We can bound  $\|\rho(t)\|_{L^2}$  easily by applying the Arnold, Falk, Winther estimate [3, p. 342], which gives

$$\|\rho(t)\|_{L^2} \leq \begin{cases} ch \|f(t)\|_{L^2} & \text{if } r = 0 \\ ch^{2+s} \|f(t)\|_{H^s} & \text{if } s \leq r - 1, \text{ for } r > 0 \end{cases} \quad (5.12)$$

Bounding  $\|\theta(t)\|_{L^2}$  is more subtle. Set  $\phi_h := \theta$  and  $\omega_h := \varepsilon$  in (5.6). Adding the equations yields

$$\frac{1}{2} \frac{d}{dt} \|\theta\|^2 + \|\varepsilon\|^2 = -(\rho_t, \theta), \quad t \in I$$

We use a technique from Thomée [27, p. 8] to derive an estimate for  $\|\theta(t)\|$ . Since  $\|\theta\|$  may not be differentiable when  $\theta = 0$ , introduce a constant  $\delta > 0$  and observe that

$$(\|\theta\|^2 + \delta^2)^{1/2} \frac{d}{dt} (\|\theta\|^2 + \delta^2)^{1/2} = \frac{1}{2} \frac{d}{dt} (\|\theta\|^2 + \delta^2) = \frac{1}{2} \frac{d}{dt} \|\theta\|^2 \leq \|\rho_t\| \|\theta\|,$$

the last step following by Cauchy-Schwarz. Since  $\|\theta\| \leq (\|\theta\|^2 + \delta^2)^{1/2}$ , we have that

$$\frac{d}{dt} (\|\theta\|^2 + \delta^2)^{1/2} \leq \|\rho_t\|.$$

Note that  $\theta(0) = u_h(0) - \tilde{u}_h(0) = g_h - g_h = 0$ . Thus

$$\|\theta(t)\| = \lim_{\delta \rightarrow 0} \int_0^t \frac{d}{dt} (\|\theta\| + \delta^2)^{1/2} \leq \int_0^t \|\rho_t\|.$$

Using the bounds on  $\|\rho(t)\|$  from (5.12), we get

$$\|\theta(t)\| \leq \begin{cases} ch \int_0^t \|f_t(\ell)\|_{L^2} d\ell & \text{if } r = 0, \\ ch^{2+s} \int_0^t \|f_t(\ell)\|_{H^s} d\ell & \text{for } r > 0, \text{ if } s \leq r - 1. \end{cases} \quad (5.13)$$

We can now assemble estimate (5.8) by collecting our results. We show the technique of the case  $r = 0$  as the other case employs identical analysis.

$$\begin{aligned} \|u_h - u\|_{L^2 \mathbb{R}^n} &= \left( \int_0^T \|u_h(t) - u(t)\|^2 dt \right)^{1/2} \\ &\leq \left( \int_0^T (\|\rho(t)\| + \|\theta(t)\|)^2 dt \right)^{1/2} \\ &\leq ch \left( \int_0^T \left( \|f(t)\| + \int_0^t \|f_t(\ell)\|_{L^2} d\ell \right)^2 dt \right)^{1/2} \\ &\leq ch \left( \int_0^T 2 \left( \|f(t)\|^2 + \left( \int_0^t \|f_t(\ell)\|_{L^2} d\ell \right)^2 \right) dt \right)^{1/2}. \end{aligned}$$

Roll the 2 into the constant  $c$  and observe that the inner integral is maximal when  $t = T$ . Thus,

$$\begin{aligned} \|u_h - u\|_{L^2 \mathbb{R}^n} &\leq ch \left( \int_0^T \|f(t)\|^2 + \|f_t\|_{L^1(I, L^2)}^2 dt \right)^{1/2} \\ &= ch \left( \|f\|_{L^2(I, L^2)}^2 + T \|f_t\|_{L^1(I, L^2)}^2 \right)^{1/2} \\ &\leq ch \left( \|f\|_{L^2(I, L^2)}^2 + T \|f_t\|_{L^1(I, L^2)}^2 + 2 \|f\|_{L^2(I, L^2)} \sqrt{T} \|f_t\|_{L^1(I, L^2)} \right)^{1/2} \\ &= ch \left( \|f\|_{L^2(I, L^2)} + \sqrt{T} \|f_t\|_{L^1(I, L^2)} \right). \end{aligned}$$

We now turn to (5.9), i.e. an error bound for the approximation of  $\sigma$ . We use the same technique of bounding  $\|\sigma(t) - \tilde{\sigma}_h(t)\|$  by the corresponding Arnold, Falk, Winther estimate and  $\|\tilde{\sigma}_h(t) - \sigma_h(t)\| (= \|\varepsilon(t)\|)$  by a modification of (5.6). First, observe that the Arnold, Falk, Winther estimate [3, p. 342] gives

$$\|\sigma(t) - \tilde{\sigma}_h(t)\| \leq ch^{1+s} \|f(t)\|_{H^s}, \text{ if } \begin{cases} s \leq r + 1, & \Lambda_h^{n-1} = \mathcal{P}_{r+1} \Lambda^{n-1}(\mathcal{T}) \\ s \leq r, & \Lambda_h^{n-1} = \mathcal{P}_{r+1}^- \Lambda^{n-1}(\mathcal{T}) \end{cases}. \quad (5.14)$$

To bound  $\|\varepsilon(t)\|$ , differentiate the second equation of (5.6) with respect to  $t$  and set  $\phi_h := 2\theta_t, \omega_h := 2\varepsilon$  yielding

$$\begin{aligned} (\theta_t, 2\theta_t) - (\operatorname{div} \varepsilon, 2\theta_t) &= -(\rho_t, 2\theta_t), \\ (\varepsilon_t, 2\varepsilon) + (\theta_t, \operatorname{div} \varepsilon) &= 0. \end{aligned}$$

Adding the equations and converting to norms, we have the bound

$$\frac{d}{dt} \|\varepsilon\|^2 + 2 \|\theta_t\|^2 = -2(\rho_t, \theta_t) \leq \|\rho_t\|^2 + \|\theta_t\|^2,$$

by Cauchy-Schwarz and the AM-GM inequality. Note that since  $\theta(0) = 0$ , we have  $\varepsilon(0) = 0$  by the second equation of (5.6). Thus

$$\|\varepsilon(t)\|^2 = \int_0^t \frac{d}{ds} \|\varepsilon(s)\|^2 ds \leq \int_0^t \frac{d}{ds} \|\varepsilon(s)\|^2 + \|\theta_t\|^2 ds \leq \int_0^t \|\rho_t\|^2 ds.$$

As before, we use (5.12) to derive

$$\|\varepsilon(t)\| \leq \begin{cases} ch \left( \int_0^t \|f_t(\ell)\|_{L^2}^2 d\ell \right)^{1/2} & \text{if } r = 0, \\ ch^{2+s} \left( \int_0^t \|f_t(\ell)\|_{H^s}^2 d\ell \right)^{1/2} & \text{for } r > 0, \text{ if } s \leq r - 1. \end{cases} \quad (5.15)$$

We assemble estimate (5.9) by collecting these results in the same fashion we did for (5.8). In the  $r = 0$  cases, we have

$$\begin{aligned} \|\sigma_h - \sigma\|_{L^2 \mathfrak{X}^{n-1}} &= \left( \int_0^T \|\sigma_h(t) - \sigma(t)\|^2 dt \right)^{1/2} \\ &\leq \left( \int_0^T (\|\sigma(t) - \tilde{\sigma}_h(t)\| + \|\varepsilon(t)\|)^2 dt \right)^{1/2} \\ &\leq c \left( \int_0^T \left( h^{1+s} \|f(t)\|_{H^s} + h \left( \int_0^t \|f_t(\ell)\|_{L^2}^2 d\ell \right)^{1/2} \right)^2 dt \right)^{1/2} \\ &\leq ch \left( \int_0^T 2 \left( h^{2s} \|f(t)\|_{H^s}^2 + \int_0^t \|f_t(\ell)\|_{L^2}^2 d\ell \right) dt \right)^{1/2}. \end{aligned}$$

Rolling the 2 into the constant  $c$  and again noting that the inner integral is maximal when  $t = T$ , we recover the first two estimates of (5.9):

$$\begin{aligned} \|\sigma_h - \sigma\|_{L^2 \mathfrak{X}^{n-1}} &\leq ch \left( \int_0^T h^{2s} \|f(t)\|_{H^s}^2 + \|f_t\|_{L^2(I, L^2)}^2 dt \right)^{1/2} \\ &= ch \left( h^{2s} \|f\|_{L^2(I, H^s)}^2 + T \|f_t\|_{L^2(I, L^2)}^2 \right)^{1/2} \\ &\leq ch \left( h^s \|f\|_{L^2(I, H^s)} + \sqrt{T} \|f_t\|_{L^2(I, L^2)} \right). \end{aligned}$$

When  $r > 0$ , (5.14) requires  $s \leq r$  or  $s \leq r + 1$  to obtain optimal convergence rates on the first term of the right side while (5.15) requires  $s \leq r - 1$  to obtain optimal rates on the second term. Thus the hypothesis  $s \leq r - 1$  implies both (5.14) and (5.15); the last estimate of (5.9) then follows by identical analysis to the first two cases.

Finally, we turn to estimate (5.10) and follow the same technique. Since  $\text{div}$  is the exterior operator  $d$  between the spaces  $\Lambda^{n-1}$  and  $\Lambda^n$ , we have from Arnold, Falk, and Winther [3, p. 342] that

$$\|\text{div}(\sigma(t) - \tilde{\sigma}_h(t))\| \leq ch^s \|f(t)\|_{H^s}, \text{ if } s \leq r + 1. \quad (5.16)$$

To bound  $\|\text{div} \varepsilon\|$  set  $w_h := \varepsilon$  in (5.6) and take the derivative with respect to  $t$ . This yields

$$\frac{d}{dt} \|\varepsilon\|^2 + (\theta_t, \text{div} \varepsilon) = 0.$$

Note that  $\operatorname{div} \varepsilon \in \Lambda_h^n$  since the discrete spaces are chosen to satisfy the relationship  $\operatorname{div} \Lambda_h^{n-1} \subset \Lambda_h^n$ . Thus, we can set  $\phi_h := \operatorname{div} \varepsilon$  in (5.6) and substitute to get

$$\frac{d}{dt} \|\varepsilon\|^2 + \|\operatorname{div} \varepsilon\|^2 = (\rho_t, \operatorname{div} \varepsilon).$$

By Cauchy-Schwarz, we have  $\|\operatorname{div} \varepsilon\|^2 \leq \|\rho_t\| \|\operatorname{div} \varepsilon\|$  and hence

$$\|\operatorname{div} \varepsilon\| \leq \|\rho_t\|.$$

Again, we use (5.12) to get

$$\|\operatorname{div} \varepsilon(t)\| \leq \begin{cases} ch \|f_t\|_{L^2} & \text{if } r = 0 \\ ch^{2+s} \|f_t\|_{H^s} & \text{if } s \leq r - 1, \text{ for } r > 0 \end{cases}. \quad (5.17)$$

The estimate (5.10) follows by combining this with (5.16).  $\square$

## 6. A Priori ERROR ESTIMATES FOR HYPERBOLIC PROBLEMS

We now analyze hyperbolic problems using the same  $L^2$  Bochner differential form spaces  $L^2 \mathcal{X}^k$  introduced in (5.1). Observe that the wave equation (2.6) is a special case of the abstract hyperbolic problem (4.5)-(4.7). Thus by Theorem 4.2, it has a unique solution on the interval  $I = [0, T]$ . For  $\Omega \subset \mathbb{R}^n$ , define the **Bochner velocity-stress mixed weak formulation**: given  $f$ ,  $u_0$ , and  $u_1$ , find  $(\mu, \sigma) \in H\mathcal{X}^n \times H\mathcal{X}^{n-1}$  such that

$$\begin{aligned} (\mu_t, \phi) - (\operatorname{div} \sigma, \phi) &= (f, \phi), \quad \forall \phi \in H\Lambda^n, \quad t \in I, \\ (\sigma_t, \omega) + (\mu, \operatorname{div} \omega) &= 0, \quad \forall \omega \in H\Lambda^{n-1}, \quad t \in I, \\ \mu(0) &= u_1, \\ \sigma(0) &= \nabla u_0, \end{aligned} \quad (6.1)$$

where  $\mu = u_t$  as in (2.8). The **semi-discrete Bochner hyperbolic problem** is: Find  $(\mu_h, \sigma_h) \in \Lambda_h^n \times \Lambda_h^{n-1}$  such that

$$\begin{aligned} (\mu_{h,t}, \phi_h) - (\operatorname{div} \sigma_h, \phi_h) &= (f, \phi_h), \quad \forall \phi_h \in \Lambda_h^n, \quad t \in I, \\ (\sigma_{h,t}, \omega_h) + (\mu_h, \operatorname{div} \omega_h) &= 0, \quad \forall \omega_h \in \Lambda_h^{n-1}, \quad t \in I, \\ \mu_h(0) &= u_{1,h}, \\ \sigma(0) &= (\nabla u_0)_h. \end{aligned} \quad (6.2)$$

The following lemma recasts notions about properties of projection operators and discrete spaces assumed by Geveci in [12] into the language of FEEC.

**Lemma 6.1.** *Choose finite element spaces*

$$\Lambda_h^{n-1} = \left\{ \begin{array}{c} \mathcal{P}_{r+1} \Lambda^{n-1}(\mathcal{T}) \\ \text{or} \\ \mathcal{P}_{r+1}^- \Lambda^{n-1}(\mathcal{T}) \end{array} \right\}, \quad \Lambda_h^n = \mathcal{P}_{r+1}^- \Lambda^n(\mathcal{T}) \quad (= \mathcal{P}_r \Lambda^n(\mathcal{T})).$$

Then the projection operators from (3.1) satisfy

$$(\operatorname{div} \pi_h^{n-1} \omega, \phi_h) = (\operatorname{div} \omega, \phi_h), \quad \forall \omega \in H(\operatorname{div}), \quad \phi_h \in \Lambda_h^n, \quad (6.3)$$

$$(\pi_h^n \phi, \operatorname{div} \omega_h) = (\phi, \operatorname{div} \omega_h), \quad \forall \phi \in L^2, \quad \omega_h \in \Lambda_h^{n-1}, \quad (6.4)$$

with the approximation properties

$$\begin{aligned} \|\pi_h^{n-1}\omega - \omega\|_{L^2\Lambda^{n-1}} &\leq ch^s \|\omega\|_{H^s\Lambda^{n-1}}, \\ &\text{for } 0 \leq s \leq r+2, \text{ if } \Lambda_h^{n-1} = \mathcal{P}_{r+1}\Lambda^{n-1}(\mathcal{T}), \text{ or} \\ &\text{for } 0 \leq s \leq r+1, \text{ if } \Lambda_h^{n-1} = \mathcal{P}_{r+1}^-\Lambda^{n-1}(\mathcal{T}), \end{aligned} \quad (6.5)$$

$$\|\pi_h^n\phi - \phi\|_{L^2\Lambda^n} \leq ch^s \|\phi\|_{H^s\Lambda^n}, \text{ for } 0 \leq s \leq r+1. \quad (6.6)$$

*Proof.* We denote  $\text{div}$  by  $d$  since it is the exterior derivative operator in this context. By Theorem 3.1 (1),  $\pi_h^{n-1}$  and  $\pi_h^n$  are projections onto Hilbert spaces and hence self-adjoint. By Theorem 3.1 (3), we have that

$$(d\pi_h^{n-1}\omega, \phi_h) = (\pi_h^n d\omega, \phi_h) = (d\omega, \pi_h^n \phi_h) = (d\omega, \phi_h),$$

establishing (6.3). A similar trick establishes (6.4). Estimate (6.5) follows directly from Theorem 3.1 (1). (Note that Theorem 3.1 (1) is stated for the case  $\mathcal{P}_r\Lambda^{n-1}$  while here we have  $\mathcal{P}_{r+1}\Lambda^{n-1}$ , thereby allowing for the higher bound on  $s$  in this case.) Finally, since  $\pi_h^n\mu - \mu = \partial_t(\pi_h^n u - u)$ , Theorem 3.1 (1) also implies (6.6).  $\square$

**Theorem 6.2.** Fix  $I := [0, T]$ . Choose finite element spaces

$$\Lambda_h^{n-1} = \left\{ \begin{array}{c} \mathcal{P}_{r+1}\Lambda^{n-1}(\mathcal{T}) \\ \text{or} \\ \mathcal{P}_{r+1}^-\Lambda^{n-1}(\mathcal{T}) \end{array} \right\}, \quad \Lambda_h^n = \mathcal{P}_{r+1}^-\Lambda^n(\mathcal{T}) \quad (= \mathcal{P}_r\Lambda^n(\mathcal{T})).$$

Then for  $(\mu_h, \sigma_h)$  the solution to (6.2), the following error estimate holds:

$$\|\mu_h - \mu\|_{L^2\mathfrak{X}^n} + \|\sigma_h - \sigma\|_{L^2\mathfrak{X}^{n-1}} \leq c \left( \sqrt{T}E_1 + h^s(\sqrt{T}E_2 + E_3 + \sqrt{T}E_4) \right), \quad (6.7)$$

where

$$E_1 = \|u_1 - u_{1,h}\|_{L^2} + \|(\nabla u_0) - (\nabla u_0)_h\|_{L^2} \quad (\text{error due to discretization of initial data})$$

$$E_2 = \|u_1\|_{H^s} + \|\nabla u_0\|_{H^s} \quad (\text{regularity of initial data})$$

$$E_3 = \|u_t\|_{L^2(I, H^s)} + \|\sigma\|_{L^2(I, H^s)} \quad (\text{regularity of continuous solution to (6.1)})$$

$$E_4 = \|u_{tt}\|_{L^2(I, H^s)} + \|\sigma_t\|_{L^2(I, H^s)} \quad (\partial_t \text{ of } E_3 \text{ accumulated over } I).$$

*Proof.* We adapt the proof technique from Geveci [12] to the FEEC setting. Define  $\Psi := \Lambda^n \times \Lambda^{n-1}$  with finite dimensional subspace  $\Psi_h := \Lambda_h^n \times \Lambda_h^{n-1}$ . Denote the components of an element  $\xi_i \in \Psi$  by  $\{\phi_i, \omega_i\}$ . The  $L^2$  inner product and norm on  $\Psi$  are

$$(\xi_1, \xi_2)_\Psi := (\phi_1, \phi_2)_{L^2} + (\omega_1, \omega_2)_{L^2} \quad \text{and} \quad \|\xi\|_\Psi := \sqrt{(\xi, \xi)_\Psi}.$$

Define a skew-symmetric bilinear form  $a : \Psi \times \Psi \rightarrow \mathbb{R}$  by

$$a(\xi_1, \xi_2) := -(\text{div } \omega_1, \phi_2)_{L^2} + (\phi_1, \text{div } \omega_2)_{L^2}.$$

Let  $\psi := (\mu, \sigma) \in \Psi$  be the solution to (6.1) and let  $\xi := (\phi, \omega) \in \Psi$  be arbitrary. Then adding the equations of (6.1) yields

$$(\psi_t, \xi)_\Psi + a(\psi, \xi) = (f, \phi)_{L^2} \quad \forall \xi \in \Psi. \quad (6.8)$$

Similarly, from (6.2) we get

$$(\psi_{h,t}, \xi_h)_\Psi + a(\psi_h, \xi_h) = (f, \phi_h)_{L^2} \quad \forall \xi_h \in \Psi_h. \quad (6.9)$$

Define a projection operator  $\pi_h : \Psi \rightarrow \Psi_h$  using the bounded cochain projections from (3.1) via

$$\pi_h \xi := (\pi_h^n \phi, \pi_h^{n-1} \omega).$$

By Lemma 6.1, we have that

$$a(\pi_h \psi, \xi_h) = a(\psi, \xi_h) \quad \forall \xi_h \in \Psi_h.$$

Further, since  $\pi_h$  only affects the spatial variables, it commutes with the time derivative operator, i.e.

$$(\partial_t \pi_h \psi, \xi_h)_\Psi = (\pi_h \partial_t \psi, \xi_h)_\Psi,$$

Combining these identities with (6.8), we have that

$$(\partial_t \pi_h \psi(t), \xi_h)_\Psi + a(\pi_h \psi(t), \xi_h) = (f(t), \phi_h)_{L^2} + (\pi_h \partial_t \psi(t) - \partial_t \psi(t), \xi_h)_\Psi, \quad \forall \xi_h \in \Psi_h \quad (6.10)$$

Now define the error function

$$\varepsilon_h(t) := \pi_h \psi(t) - \psi_h(t).$$

Subtracting (6.9) from (6.10) yields

$$(\partial_t \varepsilon_h(t), \xi_h)_\Psi + a(\varepsilon_h(t), \xi_h) = (\pi_h \partial_t \phi(t) - \partial_t \phi(t), \xi_h)_\Psi, \quad \forall \xi_h \in \Psi_h. \quad (6.11)$$

Define an operator  $\Xi_h : \Psi_h \rightarrow \Psi_h$  by

$$(\Xi_h \xi_1, \xi_2)_\Psi := a(\xi_1, \xi_2), \quad \forall \xi_1, \xi_2 \in \Psi_h.$$

Letting  $P_h : \Psi \rightarrow \Psi_h$  denote projection with respect to  $(\cdot, \cdot)_\Psi$ , we can re-write (6.11) as

$$\partial_t \varepsilon_h(t) + \Xi_h \varepsilon_h(t) = P_h (\pi_h \partial_t \phi(t) - \partial_t \phi(t)) \quad (6.12)$$

Now, since  $a$  is skew-symmetric,  $\Xi_h$  is skew-adjoint, i.e.

$$(\Xi_h \xi_1, \xi_2)_\Psi = -(\xi_1, \Xi_h \xi_2)_\Psi$$

Hence,  $-\Xi_h$  will generate the one-parameter unitary group  $e^{-t\Xi_h}$ . Therefore we can turn (6.12) into

$$\varepsilon_h(t) = e^{-t\Xi_h} \varepsilon_h(0) + \int_0^t e^{-(t-\tau)\Xi_h} P_h (\pi_h \partial_\tau \phi(\tau) - \partial_\tau \phi(\tau)) d\tau. \quad (6.13)$$

Since  $e^{-t\Xi_h}$  is unitary, it preserves  $\Psi$ -norm, i.e.  $\|e^{-t\Xi_h} \psi\|_\Psi = \|\psi\|_\Psi$  for all  $\psi \in \Psi$ . Thus (6.12) becomes

$$\|\varepsilon_h(t)\|_\Psi \leq \|\varepsilon_h(0)\|_\Psi + \int_0^t \|\pi_h \partial_\tau \phi(\tau) - \partial_\tau \phi(\tau)\|_\Psi d\tau. \quad (6.14)$$

Using (6.5) and (6.6) from Lemma 6.1, we start building up the main result. We use the notation  $f \lesssim g$  to mean  $f \leq cg$  where  $c$  is some constant independent of  $h$  and  $T$ .

$$\begin{aligned}
& (\|\mu_h - \mu\|_{L^2 \mathfrak{X}^n} + \|\sigma_h - \sigma\|_{L^2 \mathfrak{X}^{n-1}})^2 \\
& \leq 2 \int_0^T \|\mu_h - \mu\|_{L^2}^2 + \|\sigma_h - \sigma\|_{L^2}^2 dt \\
& \lesssim \int_0^T (\|\mu_h - \pi_h^n \mu\|_{L^2} + \|\sigma_h - \pi_h^{n-1} \sigma\|_{L^2})^2 \\
& \quad + \|\pi_h^n \mu - \mu\|_{L^2}^2 + \|\pi_h^{n-1} \sigma - \sigma\|_{L^2}^2 dt \\
& = \int_0^T \|\varepsilon_h(t)\|_{L^2}^2 dt + \|\pi_h^n \mu - \mu\|_{L^2 \mathfrak{X}^n}^2 + \|\pi_h^{n-1} \sigma - \sigma\|_{L^2 \mathfrak{X}^{n-1}}^2 dt \\
& \lesssim \int_0^T \|\varepsilon_h(t)\|_{L^2}^2 + h^{2s} \|\mu(t)\|_{H^s}^2 + h^{2s} \|\sigma(t)\|_{H^s}^2 dt \\
& \lesssim \int_0^T \|\varepsilon_h(t)\|_{L^2}^2 dt + h^{2s} E_3^2. \tag{6.15}
\end{aligned}$$

To bound  $\|\varepsilon_h(t)\|_{L^2}$ , we use (6.14) and the same results from Lemma 6.1 to get

$$\begin{aligned}
\|\varepsilon_h(t)\| & \lesssim \|\pi_h^n u_1 - u_{1,h}\| + \|\pi_h^{n-1}(\nabla u_0) - (\nabla u_0)_h\| \\
& \quad + \int_0^t \|\pi_h^n \partial_\tau \mu(\tau) - \partial_\tau \mu(\tau)\| + \|\pi_h^{n-1} \partial_\tau \sigma(\tau) - \partial_\tau \sigma(\tau)\| d\tau \\
& \lesssim \|u_1 - u_{1,h}\| + \|(\nabla u_0) - (\nabla u_0)_h\| + h^s (\|u_1\|_{H^s} + \|\nabla u_0\|_{H^s}) \\
& \quad + h^s \left( \int_0^t \|\partial_\tau \mu(\tau)\|_{H^s} + \|\partial_\tau \sigma(\tau)\|_{H^s} d\tau \right).
\end{aligned}$$

Integrating this from 0 to  $T$  yields

$$\begin{aligned}
\int_0^T \|\varepsilon_h(t)\|^2 dt & \lesssim T E_1^2 + h^{2s} T E_2^2 \\
& \quad + h^{2s} \int_0^T \left( \int_0^t \|\partial_\tau \mu(\tau)\|_{H^s} + \|\partial_\tau \sigma(\tau)\|_{H^s} d\tau \right)^2 dt \\
& \lesssim T E_1 + h^{2s} T E_2^2 + h^{2s} \int_0^T \left( \|\partial_\tau \mu\|_{L^2(I, H^s)} + \|\partial_\tau \sigma\|_{L^2(I, H^s)} \right)^2 dt \\
& \lesssim T E_1 + h^{2s} T E_2^2 + h^{2s} T E_4^2 \tag{6.16}
\end{aligned}$$

Combining (6.15) and (6.16) yields

$$\begin{aligned}
(\|\mu_h - \mu\|_{L^2 \mathfrak{X}^n} + \|\sigma_h - \sigma\|_{L^2 \mathfrak{X}^{n-1}})^2 & \leq c (T E_1^2 + h^{2s} T E_2^2 + h^{2s} E_3^2 + h^{2s} T E_4^2) \\
& \leq c \left( \sqrt{T} E_1 + h^s \left( \sqrt{T} E_2 + E_3 + \sqrt{T} E_4 \right) \right)^2
\end{aligned}$$

Taking the square root of both sides completes the proof.  $\square$

Theorem 6.2 implies that the error in  $u_t$  (i.e.  $\mu$ ) and the error in  $\sigma$  due to discretization of the continuous problem is bounded *a priori* over  $[0, T]$  by terms depending only on the regularity of the initial data and the continuous solution. Since the interval is finite, this implies control over the error in  $u$  as well. These results imply that the finite element choices allowed by FEEC theory for elliptic problems also result in stable semi-discrete finite element methods for hyperbolic problems.

## 7. SEMI-LINEAR EVOLUTION PROBLEMS

We now show how the techniques developed above can be extended to certain types of non-linear evolution problems. Consider the **semi-linear heat equation**: Find  $u(x, t)$  such that

$$\begin{aligned} u_t - \Delta u + F(u) &= f \quad \text{in } \Omega, \quad \text{for } t > 0 \\ u &= 0 \quad \text{on } \partial\Omega, \quad \text{for } t > 0 \quad \text{with } u(\cdot, 0) = g \text{ in } \Omega, \end{aligned} \quad (7.1)$$

where  $F$  is some non-linear operator on  $L^2(\Omega)$ . The existence and uniqueness of solutions to instances of this problem have been studied extensively [15, 13, 24, 30] as have finite element methods for the approximation of its solution [28, 9, 25, 27, 19].

We focus here on the case where  $F$  satisfies a Lipschitz condition

$$\|F(v) - F(w)\|_{L^2} \leq C \|v - w\|_{L^2}, \quad \forall v, w \in L^2(\Omega). \quad (7.2)$$

This condition, or a weaker locally Lipschitz condition [5], is assumed by Holst and Stern [17] in their recent extension of the FEEC error estimates to semi-linear elliptic problems. Hence, it serves as an obvious assumption for extending our evolution results to the semi-linear case.

For  $\Omega \subset \mathbb{R}^n$ , define the **Bochner semi-linear mixed weak form parabolic**: Given  $f$  and  $g$ , find  $(u, \sigma) \in H\mathcal{X}^n \times H\mathcal{X}^{n-1}$  such that

$$\begin{aligned} (u_t, \phi) - (\operatorname{div} \sigma, \phi) + (F(u), \phi) &= (f, \phi), \quad \forall \phi \in H\Lambda^n, \quad t \in I, \\ (\sigma, \omega) + (u, \operatorname{div} \omega) &= 0, \quad \forall \omega \in H\Lambda^{n-1}, \quad t \in I, \\ u(0) &= g. \end{aligned} \quad (7.3)$$

The **semi-linear semi-discrete Bochner parabolic problem** is thus: Find  $(u_h, \sigma_h) \in \Lambda_h^n \times \Lambda_h^{n-1}$  such that

$$\begin{aligned} (u_{h,t}, \phi_h) - (\operatorname{div} \sigma_h, \phi_h) + (F(u_h), \phi_h) &= (f, \phi_h), \quad \forall \phi_h \in \Lambda_h^n, \quad t \in I, \\ (\sigma_h, \omega_h) + (u_h, \operatorname{div} \omega_h) &= 0, \quad \forall \omega_h \in \Lambda_h^{n-1}, \quad t \in I, \\ u_h(0) &= g_h, \end{aligned} \quad (7.4)$$

where  $g_h \in \Lambda_h^n$  is an approximation of  $g$ . Analogously to the linear case, for any  $t_0 \in I$ , define the **time-ignorant linear discrete elliptic problem**: find  $(\tilde{u}_h, \tilde{\sigma}_h) \in \Lambda_h^n \times \Lambda_h^{n-1}$  such that

$$\begin{aligned} (\operatorname{div} \tilde{\sigma}_h, \phi_h) - (\Delta u(t_0), \phi_h) &= 0, \quad \forall \phi_h \in \Lambda_h^n, \quad t \in I, \\ (\tilde{\sigma}_h, \omega_h) + (\tilde{u}_h, \operatorname{div} \omega_h) &= 0, \quad \forall \omega_h \in \Lambda_h^{n-1}, \quad t \in I, \\ \tilde{u}_h(0) &= g_h, \end{aligned} \quad (7.5)$$

where now  $u$  is the solution to the continuous semi-linear problem (7.3). Similarly, define

$$\begin{aligned} \rho(t) &:= \tilde{u}_h(t) - u(t), \\ \theta(t) &:= u_h(t) - \tilde{u}_h(t), \\ \varepsilon(t) &:= \sigma_h(t) - \tilde{\sigma}_h(t), \end{aligned}$$

where  $u, \sigma$  and their discrete counterparts are now solutions to the corresponding semi-linear problems. We have an analogous lemma.

**Lemma 7.1.** *The semi-linear error functions satisfy the semi-discrete formulation, i.e.*

$$\begin{aligned} (\theta_t, \phi_h) - (\operatorname{div} \varepsilon, \phi_h) &= -(\rho_t, \phi_h), \quad \forall \phi_h \in \Lambda_h^n, \quad t \in I, \\ (\varepsilon, \omega_h) + (\theta, \operatorname{div} \omega_h) &= 0, \quad \forall \omega_h \in \Lambda_h^{n-1}, \quad t \in I. \end{aligned} \quad (7.6)$$

*Proof.* The second equation is immediate from the second equations in (7.4) and (7.5). The first equation can be written out as

$$(u_{h,t}, \phi_h) - (\operatorname{div} \sigma_h, \phi_h) + (\operatorname{div} \tilde{\sigma}_h, \phi_h) - (\tilde{u}_{h,t}, \phi_h) = (u_t, \phi_h) - (\tilde{u}_{h,t}, \phi_h)$$

which is reduced as follows:

$$\begin{aligned} (u_{h,t}, \phi_h) - (\operatorname{div} \sigma_h, \phi_h) + (\operatorname{div} \tilde{\sigma}_h, \phi_h) &= (u_t, \phi_h) && \text{cancel like terms} \\ (u_{h,t}, \phi_h) - (\operatorname{div} \sigma_h, \phi_h) &= -(\Delta u, \phi_h) + (u_t, \phi_h) && \text{by (7.5)} \\ (f, \phi_h) - (F(u_h), \phi_h) &= -(\Delta u, \phi_h) + (u_t, \phi_h) && \text{by (7.4)} \end{aligned}$$

This says that the continuous problem  $u_t - \Delta u + F(u_h) = f$  should hold in a weak sense when tested against any of the functions in  $\Lambda_h^n$ . This is guaranteed to be true since we chose  $\Lambda_h^n \subset \Lambda^n = L^2$ . Thus, the error equations hold as stated.  $\square$

**Theorem 7.2.** Fix  $I := [0, T]$ . Suppose  $(u, \sigma)$  is the solution to (7.3) such that the regularity estimate

$$\|u(t)\|_{H^{s+2}} + \|du(t)\|_{H^{s+1}} + \|\sigma(t)\|_{H^{s+1}} + \|d\sigma(t)\|_{H^s} \leq c \|f(t)\|_{H^s} \quad (7.7)$$

holds for  $0 \leq s \leq s_{\max}$  and  $t \in I$ . Assume that the operator  $F$  satisfies the Lipschitz assumption (7.2). Choose finite element spaces

$$\Lambda_h^{n-1} = \left\{ \begin{array}{c} \mathcal{P}_{r+1} \Lambda^{n-1}(\mathcal{T}) \\ \text{or} \\ \mathcal{P}_{r+1}^- \Lambda^{n-1}(\mathcal{T}) \end{array} \right\}, \quad \Lambda_h^n = \mathcal{P}_{r+1}^- \Lambda^n(\mathcal{T}) \quad (= \mathcal{P}_r \Lambda^n(\mathcal{T}))$$

Then for  $0 \leq s \leq s_{\max}$ ,  $g_h$  defined by (5.4), and  $(u_h, \sigma_h)$  the solution to (7.4), the following error estimates hold:

$$\|u_h - u\|_{L^2 \mathfrak{X}^n} \leq \begin{cases} ch \left( \|f\|_{L^2(I, L^2)} + \sqrt{T} \|f_t\|_{L^1(I, L^2)} \right) & \text{if } r = 0 \\ ch^{1+s} \left( \|f\|_{L^2(I, H^s)} + \sqrt{T} \|f_t\|_{L^1(I, H^s)} \right) & \text{for } r > 0, \text{ if } s \leq r - 1 \end{cases} \quad (7.8)$$

$$\|\sigma_h - \sigma\|_{L^2 \mathfrak{X}^{n-1}} \leq \begin{cases} ch \left( \|f\|_{L^2(I, H^s)} + \sqrt{T} \|f_t\|_{L^2(I, L^2)} \right) \\ \quad \text{if } r = 0, s = 0, \Lambda_h^{n-1} = \mathcal{P}_1^- \Lambda^{n-1}(\mathcal{T}) \\ c \left( h^{1+s} \|f\|_{L^2(I, H^s)} + h \sqrt{T} \|f_t\|_{L^2(I, L^2)} \right) \\ \quad \text{if } r = 0, s \leq 1, \Lambda_h^{n-1} = \mathcal{P}_1 \Lambda^{n-1}(\mathcal{T}) \\ c \left( h^{1+s} \|f\|_{L^2(I, H^s)} + h^{(3/2)+s} \sqrt{T} \|f_t\|_{L^2(I, H^s)} \right) \\ \quad \text{for } r > 0, \text{ if } s \leq r - 1 \end{cases} \quad (7.9)$$

$$\|\operatorname{div}(\sigma_h - \sigma)\|_{L^2 \mathfrak{X}^n} \leq \begin{cases} c \left( h^s \|f\|_{L^2(I, H^s)} + h \|f_t\|_{L^2(I, L^2)} \right) \\ \quad \text{if } r = 0, s \leq 1 \\ c \left( h^s \|f\|_{L^2(I, H^s)} + h^{2+s} \|f_t\|_{L^2(I, H^s)} \right) \\ \quad \text{for } r > 0, \text{ if } s \leq r - 1 \end{cases} \quad (7.10)$$

*Proof.* The proof is very similar to that of Theorem 5.2. Equation (7.5) is the  $k = n$  case of the discrete mixed variational problem examined by Holst and Stern in [17, Equation (9)]. Therefore, we can use the same type of triangle inequality from (5.11) to recover the estimates. By [17, Theorem 4.2], we have the estimates

$$\|\rho(t)\|_{L^2} \leq \begin{cases} ch\|f(t)\|_{L^2} & \text{if } r = 0 \\ ch^{2+s}\|f(t)\|_{H^s} & \text{if } s \leq r - 1, \text{ for } r > 0 \end{cases} \quad (7.11)$$

$$\|\sigma(t) - \tilde{\sigma}_h(t)\|_{L^2} \leq ch^{1+s}\|f(t)\|_{H^s}, \text{ if } \begin{cases} s \leq r + 1, & \Lambda_h^{n-1} = \mathcal{P}_{r+1}\Lambda^{n-1}(\mathcal{T}) \\ s \leq r, & \Lambda_h^{n-1} = \mathcal{P}_{r+1}^-\Lambda^{n-1}(\mathcal{T}) \end{cases} \quad (7.12)$$

$$\|\operatorname{div}(\sigma(t) - \tilde{\sigma}_h(t))\|_{L^2} \leq ch^s\|f(t)\|_{H^s}, \text{ if } s \leq r + 1. \quad (7.13)$$

An explanation of how these estimates are derived from the results of [17] is given in Appendix A. Note that these estimates are exactly the same as the corresponding estimates (5.12), (5.14) and (5.16) from the linear case. The proof then proceeds exactly as before since the rest of the argument does not appeal to the linearity of the problem at all.  $\square$

This approach seems likely to extend to semi-linear hyperbolic problems as well. Since the well-posedness of such problems is a significant issue in its own right, however, we do not consider such an approach in the present work.

## 8. CONCLUDING REMARKS

In this article, we have extended the Finite Element Exterior Calculus of Arnold, Falk, and Winther [3, 2] for linear mixed variational problems to linear and semi-linear parabolic and hyperbolic evolution systems. Both the parabolic and hyperbolic cases make strong use of the smoothed projection operators  $\pi_h^k$ , which are one of the most elaborate and delicate constructions in the FEEC framework. In the parabolic case, the use of the  $\pi_h^k$  operators was hidden somewhat by the use of elliptic projection error estimates, proofs of which rely on properties of these operators. In the hyperbolic case, the proof techniques use these properties more explicitly. In any case, the formal treatment and generalization of these operators by Arnold, Falk and Winther can now be seen as a useful tool for the analysis of evolution problems as well as elliptic PDE.

We have also seen in this article how the recent generalizations of the FEEC by Holst and Stern [16, 17] for semi-linear elliptic PDE can be extended to evolution PDE as well, both parabolic and hyperbolic types. We also anticipate that the basic approach to analyzing variational crimes in [16, 17] for the linear and semilinear elliptic cases will also work in the case of evolution problems; we will explore the question of variational crimes in a subsequent article, with the target being the analysis of surface finite element methods for evolution problems.

## 9. ACKNOWLEDGMENTS

MH was supported in part by NSF Awards 0715146, by DOD/DTRA Award HDTRA-09-1-0036, and by NBCR. AG was supported in part by NSF Award 0715146 and by NBCR.

## APPENDIX A. EXPLANATION OF SEMI-LINEAR ERROR ESTIMATES

In this appendix, we explain why estimates (7.11), (7.12), and (7.13) follow from [17, Theorem 4.2]. We will focus just on the  $r > 0$  case of (7.11) as it requires the sharpening

of a special case of an estimate appearing in [17, Theorem 4.2]. The other cases work out along similar lines by a direct application of the Holst and Stern estimates.

First, we recall some notation from [3] used in [17]. If  $(W, d)$  is a Hilbert complex with associated domain complex  $(V, d)$  and parametrized subcomplex family  $(V_h, d)$ , denote the best approximation in  $W$ -norm by

$$E(w) = \inf_{v \in V_h^k} \|w - v\|_W, \quad w \in W^k.$$

The relevant result from Holst and Stern [17, Theorem 4.2] is stated as

$$\begin{aligned} \|u - \tilde{u}_h\|_V + \|p - p_h\|_W &\leq c(E(u) + E(du) + E(p)) \\ &\quad + \eta[E(\sigma) + E(d\sigma)] + (\delta + \mu)E(d\sigma) + \mu E(P_{\mathfrak{B}}u), \end{aligned} \tag{A.1}$$

where  $\eta$ ,  $\delta$ , and  $\mu$  are coefficients defined as the norms of certain abstract operators,  $u \in W_k$ , and  $p$  is a harmonic  $k$ -form with discrete counterpart  $p_h$  introduced to make the abstract Hodge-Laplacian problem well-posed.

Casting this into the context of the deRham complex, we have

$$(W, d) = (L^2\Lambda, d) \quad \text{and} \quad (V, d) = (H\Lambda, d).$$

Since we are interested here only in the case  $k = n$ , there are no harmonic  $k$ -forms so that  $p = p_h = 0$ . Further,  $du = 0$  since  $d\Lambda^n = 0$ , whereby  $\|u - \tilde{u}_h\|_V = \|u - \tilde{u}_h\|_W = \|u - \tilde{u}_h\|_{L^2}$ . This eliminates the error terms in  $p$  and  $du$ , giving us the reduced estimate

$$\|u - \tilde{u}_h\|_{L^2} \leq c(E(u) + \eta[E(\sigma) + E(d\sigma)] + (\delta + \mu)E(d\sigma) + \mu E(P_{\mathfrak{B}}u)).$$

Crucially, this estimate can be reduced further when  $k = n$ . The derivation of (A.1) uses the estimate

$$\|d(u - \tilde{u}_h)\|_W \leq c(E(du) + \eta[E(d\sigma) + E(p)])$$

from [3, Theorem 2.11] which is unnecessary here since the left side is always zero. Since this is the only part of the derivation that requires the term  $\eta E(d\sigma)$ , we can drop it, yielding

$$\|u - \tilde{u}_h\|_{L^2} \leq c(E(u) + \eta E(\sigma) + (\delta + \mu)E(d\sigma) + \mu E(P_{\mathfrak{B}}u)). \tag{A.2}$$

We now give bounds on each of the terms in (A.2). The coefficients appearing in the abstract estimates can be stated in terms of powers of  $h$  in the deRham context. These appear in [3, p. 312] as

$$\eta = O(h), \quad \delta = O(h^{\min(2, r+1)}), \quad \text{and} \quad \mu = O(h^{r+1}).$$

To bound the error terms, Arnold, Falk and Winther define smooth projection operators  $\pi_h^k : L^2\Lambda^k(\Omega) \rightarrow \Lambda_h^k$  satisfying optimal convergence rates as stated precisely in [3, Theorem 5.9]. For instance, if  $\Lambda_h^k$  is one of  $\mathcal{P}_{r+1}^-\Lambda^k(\mathcal{T}_h)$  or, if  $r \geq 1$ ,  $\mathcal{P}_r\Lambda^k(\mathcal{T}_h)$  then

$$\|w - \pi_h^k w\|_{L^2\Lambda^k(\Omega)} \leq ch^s \|w\|_{H^s\Lambda^k(\Omega)}, \quad \text{for } w \in H^s\Lambda^k(\Omega), \quad 0 \leq s \leq r+1.$$

These types of results bound  $E(w)$  in terms of  $\|w\|_{H^s\Lambda^k}$ , which is in turn bounded in terms of  $\|f\|_{H^s\Lambda^k}$  by the regularity hypothesis (7.7). Summarizing these results, we have

$$\begin{aligned} E(u) &\leq ch^{s+2} \|f\|_{H^s} \\ E(\sigma) &\leq ch^{s+1} \|f\|_{H^s} \\ E(d\sigma) &\leq ch^s \|f\|_{H^s} \\ E(P_{\mathfrak{B}}u) &\leq ch^{s+2} \|f\|_{H^s} \end{aligned}$$

We can now prove (7.11) by collecting results and applying them to (A.2), yielding

$$\|\rho(t)\|_{L^2} \leq c(h^{s+2} + h(h^{s+1}) + (h^{\min(2,r+1)} + h^{r+1})h^s + h^{r+1}h^{s+2}) \|f(t)\|_{H^s}$$

The greatest common factor from the above expression is  $h^{s+2}$  hence this is the overall order estimate that can be inferred, as was claimed.

## REFERENCES

- [1] D. Arnold and H. Chen. Finite element exterior calculus for parabolic equations. *In preparation*, 2012.
- [2] D. Arnold, R. Falk, and R. Winther. Finite element exterior calculus, homological techniques, and applications. *Acta Numerica*, pages 1–155, 2006.
- [3] D. Arnold, R. Falk, and R. Winther. Finite element exterior calculus: from Hodge theory to numerical stability. *Bulletin of the American Mathematical Society*, 47(2):281–354, 2010.
- [4] G. Baker. Error estimates for finite element methods for second order hyperbolic equations. *SIAM journal on numerical analysis*, pages 564–576, 1976.
- [5] R. E. Bank, M. J. Holst, R. Szypowski, and Y. Zhu. Finite element error estimates for critical growth semilinear problems without angle conditions. *arXiv:1108.3661*, 2011.
- [6] A. Bossavit. Whitney forms: a class of finite elements for three-dimensional computations in electromagnetism. *Science, Measurement and Technology, IEE Proceedings A*, 135(8):493–500, nov 1988.
- [7] J. Bramble, A. Schatz, V. Thomée, and L. Wahlbin. Some convergence estimates for semidiscrete galerkin type approximations for parabolic equations. *SIAM Journal on Numerical Analysis*, pages 218–241, 1977.
- [8] J. Brüning and M. Lesch. Hilbert complexes. *J. Funct. Anal.*, 108(1):88–132, 1992.
- [9] K. Chrysafinos and L. Hou. Error estimates for semidiscrete finite element approximations of linear and semilinear parabolic equations under minimal regularity assumptions. *SIAM journal on numerical analysis*, pages 282–306, 2003.
- [10] L. Cowsar, T. Dupont, and M. Wheeler. A priori estimates for mixed finite element methods for the wave equation. *Computer Methods in Applied Mechanics and Engineering*, 82(1-3):205–222, 1990.
- [11] L. Cowsar, T. Dupont, and M. Wheeler. A priori estimates for mixed finite element approximations of second-order hyperbolic equations with absorbing boundary conditions. *SIAM journal on numerical analysis*, pages 492–504, 1996.
- [12] T. Geveci. On the application of mixed finite element methods to the wave equations. *Mathematical Modelling and Numerical Analysis*, 22(2):243–250, 1988.
- [13] Y. Giga. A bound for global solutions of semilinear heat equations. *Communications in mathematical physics*, 103(3):415–421, 1986.
- [14] P. W. Gross and P. R. Kotiuga. *Electromagnetic theory and computation: a topological approach*, volume 48 of *Mathematical Sciences Research Institute Publications*. Cambridge University Press, Cambridge, 2004.
- [15] D. Henry and S. S. en ligne). *Geometric theory of semilinear parabolic equations*, volume 61. Springer-Verlag Berlin, 1981.
- [16] M. Holst and A. Stern. Geometric variational crimes: Hilbert complexes, finite element exterior calculus, and problems on hypersurfaces. *Arxiv preprint arXiv:1005.4455*, 2010.
- [17] M. Holst and A. Stern. Semilinear mixed problems on hilbert complexes and their numerical approximation. *Found. Comput. Math.*, in press, 2010.
- [18] C. Johnson and V. Thomee. Error estimates for some mixed finite element methods for parabolic type problems. *RAIRO Anal. Numér.*, 15(1):41–78, 1981.
- [19] A. Lasis and E. Süli. *hp*-version discontinuous galerkin finite element method for semilinear parabolic problems. *SIAM Journal on Numerical Analysis*, 45(4):1544–1569, 2007.
- [20] J.-C. Nédélec. Mixed finite elements in  $\mathbb{R}^3$ . *Numer. Math.*, 35(3):315–341, 1980.
- [21] J.-C. Nédélec. A new family of mixed finite elements in  $\mathbb{R}^3$ . *Numer. Math.*, 50(1):57–81, 1986.
- [22] P.-A. Raviart and J. M. Thomas. A mixed finite element method for 2nd order elliptic problems. In *Mathematical aspects of finite element methods (Proc. Conf., Consiglio Naz. delle Ricerche (C.N.R.), Rome, 1975)*, pages 292–315. Lecture Notes in Math., Vol. 606. Springer, Berlin, 1977.
- [23] M. Renardy and R. Rogers. *An introduction to partial differential equations*, volume 13. Springer Verlag, 2nd edition, 2004.

- [24] B. Ruf and E. Terraneo. The cauchy problem for a semilinear heat equation with singular initial data. In *Evolution equations, semigroups and functional analysis: in memory of Brunello Terreni*, volume 50. Birkhauser, 2002.
- [25] T. Sun and D. Filippova. Long-time error estimation for semi-linear parabolic equations. *Journal of Computational and Applied Mathematics*, 185(1):1 – 18, 2006.
- [26] R. Temam. *Infinite-dimensional dynamical systems in mechanics and physics*, volume 68. Springer verlag, 1997.
- [27] V. Thomée. *Galerkin finite element methods for parabolic problems*. Springer Verlag, 2006.
- [28] V. Thomee and L. Wahlbin. On galerkin methods in semilinear parabolic problems. *SIAM Journal on Numerical Analysis*, pages 378–389, 1975.
- [29] M. Wheeler. A priori  $L^2$  error estimates for galerkin approximations to parabolic partial differential equations. *SIAM Journal on Numerical Analysis*, pages 723–759, 1973.
- [30] J. Wrkich and Q. Zhang. Solvability conditions for some semi-linear parabolic equations. *Journal of Differential Equations*, 247(8):2440–2454, 2009.

*E-mail address:* akgillette@ucsd.edu

*E-mail address:* mholst@math.ucsd.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA SAN DIEGO, LA JOLLA CA 92093